C^*-ALGEBRAS ASSOCIATED WITH TOPOLOGICAL GROUP QUIVERS I:
GENERATORS, RELATIONS AND SPATIAL STRUCTURE

SHAWN J. M$^{\mathrm{c}}$CANN

arXiv:1212.3397v2 [math.OA] 1 Feb 2013

Abstract. Topological quivers generalize the notion of directed graphs in which the sets of vertices and edges are locally compact (second countable) Hausdorff spaces. Associated to a topological quiver Q is a C^*-correspondence, and in turn, a Cuntz-Pimsner algebra $C^*(Q)$. Given Γ a locally compact group and α and β endomorphisms on Γ, one may construct a topological quiver $Q_{\alpha,\beta}(\Gamma)$ with vertex set Γ, and edge set $\Omega_{\alpha,\beta}(\Gamma) = \{(x,y) \in \Gamma \times \Gamma \,|\, \alpha(y) = \beta(x)\}$. In this paper, the author examines the Cuntz-Pimsner algebra $\mathcal{O}_{\alpha,\beta}(\Gamma) := C^*(Q_{\alpha,\beta}(\Gamma))$. The investigative topics include a notion for topological quiver isomorphisms, generators (and their relations) of the C^*-algebras $\mathcal{O}_{\alpha,\beta}(\Gamma)$, and its spatial structure (i.e., colimits, tensor products and crossed products) and a few properties of its C^*-subalgebras.

1. Introduction and Notation

1.1. **Background.** The study of directed graphs have played a pivotal role in the construction and analysis of C^*-algebras ultimately describing certain C^*-algebras in terms of easily manageable and computatable relations. That is, given a directed graph $G = (V, E, r, s)$ with vertex set V, edge set E, range map r and source map s, one may produce a (universal) C^*-algebra generated by projections $\{p_v\}_{v\in V}$ and partial isometries $\{s_e\}_{e\in E}$ satisfying the relations

(1) $s_e^* s_e = p_{r(e)}$ for all $e \in E$
(2) $p_x = \sum_{s(e)=x} s_e s_e^*$ for all $x \in X$ with $0 < |s^{-1}(x)| < \infty$
(3) $s_e s_e^* \leq p_{s(e)}$ for all $e \in E$.

Many C^*-algebras, such as Cuntz-algebras [14], Cuntz-Krieger algebras [15], the Toeplitz algebra and the n by n matrices, are able to be defined by generators and relations which correspond directly to such a graph. Other C^*-algebras, including some that arise in representation theory (see [49, 50, 58]) and the theory of quantum spaces (see [29, 30, 31]), are of considerable importance and interest in determining how they may be represented by a graph.

Schweizer ([61]) introduced continuous (directed) graphs and so began the analysis of spaces of vertices and edges that are topological spaces. Perhaps, the first to analyze prototypes of this structure was Deaconu (see [17, 18, 19, 20]) who was interested in groupoid representations of Cuntz-like C^*-algebras associated to (non-invertible) maps of topological spaces, i.e. to endomorphisms of $C_0(X)$. This led Deaconu and Muhly [21] to extend the notion to branched coverings. In order to study these C^*-algebras, and in particular to compute its K-theory, it was beneficial to use machinery pioneered by Pimsner in his seminal paper [60] and realize the C^*-algebra as what is now called a Cuntz-Pimsner algebra. Pimsner's work was profoundly influenced by the theory of certain graph C^*-algebras (in particular, Cuntz-Krieger algebras) and crossed products by $\mathbb{Z}$ independent of the study of "continuous graph C^*-algebras."

The notion of topological quivers generalizes that of directed graphs. This is a quintuple $Q = (X, E, r, s, \lambda)$, where X and E are locally compact (second countable) Hausdorff spaces, r and s are continuous maps from X to E with r open, and $\lambda = \{\lambda_x\}_{x \in E}$ is a system of Radon measures. With topological quiver, one can associate a Cuntz-Pimsner C^*-algebra $C^*(Q)$ that also generalizes the notion of graph C^*-algebra. In a purely set-theoretic perspective, Q is a directed graph and the system of measures is to enable one to replace sums (that arise in the graph theoretic setting) with integrals. The term "topological quiver" was adopted from the ring-theoretic use of the term quiver which was introduced by Gabriel in [28] to describe the graph associated to a so-called basic algebra.

Two particularly important works were the first to investigate general structures of this nature. The first is that of Katsura [38, 39, 40, 41, 42] where the term topological graph is used to denote a structure similar to a topological quiver (X, E, r, s) with the addition restriction that r is a local homeomorphism. If λ is taken to be counting measures on the fibers of r then it has been shown that the C^*-algebra in Katsura's work is isomorphic to the C^*-algebra produced by considering the topological quiver $Q = (X, E, r, s, \lambda)$. The second paper is [7] where Brenken studies topological relations (X, E, r, s, λ) where E is a closed subset of $X \times X$. The maps r and s are the projections of this set into X. Such topological relations can be viewed as multiplicity one topological quivers. The topological quivers considered in this paper are examples of topological relations.

In [24], Exel, an Huef and Raeburn define C^*-algebras associated with a system (B, α, L) where α is an endomorphism of a unital C^*-algebra B and L is a positive linear map $L : B \to B$ such that $L(\alpha(a)b) = aL(b)$ for all $a, b \in B$ called a *transfer operator*. In fact, the C^*-algebra they generate is a Cuntz-Pimsner algebra and under certain restrictions, a C^*-algebra associated with a topological quiver; in particular, when $B = C(\mathbb{T}^d)$ the continuous function on the d-torus, $F \in M_d(\mathbb{Z})$ and α is the endomorphism

$$\alpha(f)(e^{2\pi i t}) = f(e^{2\pi i F t})$$

for $f \in C(\mathbb{T}^d)$ and $t \in \mathbb{R}^d$. Furthermore, Yamashita [68] considers topological relations of the form $Q_{n,m}(\mathbb{T}) = (\mathbb{T}, \Omega_{n,m}(\mathbb{T}), r, s, \lambda)$ where $n, m \in \mathbb{Z}$, $\gcd(n, m) = 1$ and

$$\Omega_{n,m}(\mathbb{T}) = \{(x, y) \in \mathbb{T} \times \mathbb{T} \,|\, y^n = x^m\}.$$

He then presents the C^*-algebra associated with $Q_{n,m}(\mathbb{T})$ as a universal C^*-algebra with certain generators and relation; that is, the unversal C^*-algebra generated by a unitary U and isometries $S_1, ..., S_n$ such that

$$US_i = S_{i+1}\,(1 \le i \le n-1), \qquad US_n = S_1 U^m, \qquad \sum_{i=1}^{n} S_i S_i^* = 1.$$

We generalize the notions considered in [24] and [68] and provide an extensive survey.

In Section 2, the necessary background material is provided in order to define topological quivers and the C^*-algebra associated with a topological quiver. We then develop a method for determining a presentation of the universal C^*-algebra and define a notion for a topological quiver isomorphism and prove this morphism gives an isomorphism between the associated C^*-algebras. Section 3 then contains the definition of a particular topological quiver, called a topological group relation, defined using a locally compact group Γ and endomorphisms α and β; that is,

$$Q_{\alpha,\beta}(\Gamma) = (\Gamma, \Omega_{\alpha,\beta}(\Gamma), r, s, \lambda)$$

where

$$\Omega_{\alpha,\beta}(\mathbb{T}^d) = \{(x,y) \in \Gamma \times \Gamma \,|\, \alpha(y) = \beta(x)\}.$$

For instance, if $F, G \in M_d(\mathbb{Z})$ where $\det F, \det G \neq 0$, then

$$Q_{F,G}(\mathbb{T})^d = (\mathbb{T}^d, \Omega_{F,G}(\mathbb{T}^d), r, s, \lambda)$$

where

$$\Omega_{F,G}(\mathbb{T}^d) = \{(x,y) \in \mathbb{T}^d \times \mathbb{T}^d \,|\, \sigma_F(y) = \sigma_G(x)\}$$

and $\sigma_F(e^{2\pi i t}) = e^{2\pi i F t}$ for $t \in \mathbb{R}^d$. These topological group relations are a generalization of the topological quivers studied in [24] and [68] and they also share many similar properties with the Cuntz algebra. We also consider many other examples of topological group relations and also presents the C^*-algebras, $\mathcal{O}_{F,G}(\mathbb{T}^d)$ associated with $Q_{F,G}(\mathbb{T}^d)$ as a universal C^*-algebra generated by commuting unitaries $\{U_j\}_{j=1}^d$ and an isometry S satisfying certain relations:

(1) $S^*U^\nu S = \delta_\nu^{\nu'}$,
(2) $U_j^{a_j} S = SU^{G_j}$, for all $j = 1, ..., d$ and
(3) $1 = \sum_{\nu \in \mathfrak{I}(F)} U^\nu SS^* U^{-\nu}$

where U^ν denotes $\prod_{j=1}^d U_j^{\nu_j}$ and $\mathfrak{I}(F) = \{\nu = (\nu_j) \in \mathbb{Z} \,|\, 0 \leq \nu_j \leq a_j - 1\}$. In Section 4, we investigate a few properties of C^*-subalgebras of $\mathcal{O}_{F,G}(\mathbb{T}^d)$ while in Section 5, we examine the spatial structure of $\mathcal{O}_{F,G}(\mathbb{T}^d)$ and present $\mathcal{O}_{F,G}(\mathbb{T}^d)$ as the crossed product by an endomorphism of a colimit. In [62], Stacey defines the *crossed product by an endomorphism*, ρ, as the universal C^*-algebra $B \rtimes_\rho \mathbb{N}$ generated by (a copy of) B and an isometry s such that $\rho(b) = sbs^*$ for all $b \in B$. Ultimately, it is shown that

$$\mathcal{O}_{F,G}(\mathbb{T}^d) = (\mathrm{colim}_k M_{N^k}(C(\mathbb{T}^d))) \rtimes_\rho \mathbb{N}$$

where the isometry s is, none other than, S.

1.2. **Notation.** The sets of natural numbers, integers, rationals numbers, real numbers and complex numbers will be denoted by $\mathbb{N}$, $\mathbb{Z}$, $\mathbb{Q}$, $\mathbb{R}$, and $\mathbb{C}$, respectively. Convention: $\mathbb{N}$ does not contain zero. Finally, $\mathbb{Z}_p$ denotes the abelian group $\mathbb{Z}/p\mathbb{Z} = \{0, 1, ..., p-1 \mod p\}$ and $\mathbb{T}$ denotes the torus $\{z \in \mathbb{C} \,|\, |z| = 1\}$. Whenever convenient, view $\mathbb{Z}_p \subset \mathbb{T}$ by $\mathbb{Z}_p \cong \{z \in \mathbb{T} \,|\, z^p = 1\}$.

For a topological space Y, the closure of Y is denoted $\overline{Y}$. Given a locally compact Hausdorff space X, let $C(X)$ be the continuous complex functions on X, $C_0(X)$ be the continuous complex functions on X vanishing at infinity, and $C_c(X)$ be the continuous complex functions on X with compact support. The supremum norm, denoted $||\cdot||_\infty$, is defined by

$$||f||_\infty = \sup_{x\in X}\{|f(x)|\}$$

for each continuous map $f : X \to \mathbb{C}$. For a continuous function $f \in C_c(X)$, denote the open support of f by osupp $f = \{x \in X \,\big|\, f(x) \neq 0\}$ and the support of f by supp $f = \overline{\text{osupp} f}$.

For C^*-algebras A and B, A is isomorphic to B will be written $A \cong B$; for example, we use $C(\mathbb{T}^d) \otimes M_N(\mathbb{C}) \cong M_N(C(\mathbb{T}^d))$. Moreover, $A^{\oplus n}$ denotes the n-fold direct sum $A\oplus\cdots\oplus A$. Given a group Γ and a ring R, a normal subgroup, N, of Γ is denoted $N \lhd \Gamma$ and an ideal, I, of R is denoted $I \lhd R$. Note if R is a C^*-algebra then the term ideal denotes a closed two-sided ideal. Furthermore, End(Γ) (End(R)) and Aut(Γ) (Aut(R)) denotes the set of endomorphisms of Γ (R) and automorphisms of Γ (R), respectively. For a map $\gamma : \Gamma \to \text{Aut}(A)$, the *fixed point set* is denoted A^γ and defined by

$$A^\gamma = \{a \in A \,\big|\, \gamma(g)(a) = a \text{ for each } g \in \Gamma\}.$$

Let $\alpha \in C(X)$ then $\alpha^\# \in \text{End(C(X))}$ denotes the endomorphism of $C(X)$ defined by

$$\alpha^\#(f) = f \circ \alpha \qquad \text{for each } f \in C(X).$$

Let S be a set and define the Kronecker delta function $\delta : S \times S \to \{0, 1\}$ by

$$\delta_s^r := \delta(s, r) = \begin{cases} 0 & \text{if } s \neq r \\ 1 & \text{if } s = r \end{cases}.$$

2. Preliminairies

2.1. Hilbert C^*-modules.

Definition 2.1. [48] If A is a C^*-algebra, then a *(right) Hilbert A-module* is a Banach space $\mathcal{E}_A$ together with a right action of A on $\mathcal{E}_A$ and an A-valued inner product $\langle\cdot,\cdot\rangle_A$ satisfying

(1) $\langle\xi, \eta a\rangle_A = \langle\xi, \eta\rangle_A a$
(2) $\langle\xi, \eta\rangle_A = \langle\eta, \xi\rangle_A^*$
(3) $\langle\xi, \xi\rangle \geq 0$ and $||\xi|| = ||\langle\xi, \xi\rangle_A^{1/2}||_A$

for all ξ, $\eta \in \mathcal{E}_A$ and $a \in A$ (if the context is clear, we denote $\mathcal{E}_A$ simply by $\mathcal{E}$). For Hilbert A-modules $\mathcal{E}$ and $\mathcal{F}$, call a function $T : \mathcal{E} \to \mathcal{F}$ *adjointable* if there is a function $T^* : \mathcal{F} \to \mathcal{E}$ such that $\langle T(\xi), \eta\rangle_A = \langle\xi, T^*(\eta)\rangle_A$ for all $\xi \in \mathcal{E}$ and $\eta \in \mathcal{F}$. Let $\mathcal{L}(\mathcal{E}, \mathcal{F})$ denote the set of adjointable (A-linear) operators from $\mathcal{E}$ to $\mathcal{F}$. If $\mathcal{E} = \mathcal{F}$, then $\mathcal{L}(\mathcal{E}) := \mathcal{L}(\mathcal{E}, \mathcal{E})$ is a C^*-algebra. Let $\mathcal{K}(\mathcal{E}, \mathcal{F})$ denote the closed two-sided ideal of *compact operators* given by

$$\mathcal{K}(\mathcal{E}, \mathcal{F}) := \overline{\text{span}}\{\theta_{\xi,\eta}^{\mathcal{E},\mathcal{F}} \,\big|\, \xi \in \mathcal{E},\, \eta \in \mathcal{F}\}$$

where

$$\theta_{\xi,\eta}^{\mathcal{E},\mathcal{F}}(\zeta) = \xi\langle\eta, \zeta\rangle_A \qquad \text{for each } \zeta \in \mathcal{E}.$$

Similarly, $\mathcal{K}(\mathcal{E}) := \mathcal{K}(\mathcal{E},\mathcal{E})$ and $\theta^{\mathcal{E}}_{\xi,\eta}$ (or $\theta_{\xi,\eta}$ if understood) denotes $\theta^{\mathcal{E},\mathcal{E}}_{\xi,\eta}$. For Hilbert A-module $\mathcal{E}$, the linear span of $\{\langle\xi,\eta\rangle \,|\, \xi,\eta\in\mathcal{E}\}$, denoted $\langle\mathcal{E},\mathcal{E}\rangle$, once closed is a two-sided ideal of A. Note that $\mathcal{E}\langle\mathcal{E},\mathcal{E}\rangle$ is dense in $\mathcal{E}$. The Hilbert module $\mathcal{E}$ is called *full* if $\langle\mathcal{E},\mathcal{E}\rangle$ is dense in A. The Hilbert module A_A refers to the Hilbert module A over itself, where $\langle a,b\rangle = a^*b$ for all $a,b\in A$.

An *algebraic generating set* for $\mathcal{E}$ is a subset $\{u_i\}_{i\in\mathcal{I}}\subset\mathcal{E}$ for some indexing set $\mathcal{I}$ such that $\mathcal{E}$ equals the linear span of $\{u_i\cdot a \,|\, i\in\mathcal{I}, a\in A\}$.

Definition 2.2. [37] A subset $\{u_i\}_{i\in\mathcal{I}}\subset\mathcal{E}$ is called a *basis* provided the following reconstruction formula holds for all $\xi\in\mathcal{E}$:

$$\xi = \sum_{i\in\mathcal{I}} u_i\cdot\langle u_i,\xi\rangle \qquad (\text{in } \mathcal{E}, ||\cdot||.)$$

If $\langle u_i,u_j\rangle = \delta_i^j$ as well, call $\{u_i\}_{i\in\mathcal{I}}$ an *orthonormal basis* of $\mathcal{E}$.

Remark 2.3. The preceding definition is in accordance with the finite version in [37], but many other versions exist such as in [24] where $\{u_i\}_{i=1}^n$ is called a finite Parseval frame, or in [68] where this is taken as the definition for *finitely generated.* There has been substantial work done on similar frames (see [32]).

Definition 2.4. [10, 11] If A and B are C^*-algebras, then an $A{-}B$ *C^*-correspondence* $\mathcal{E}$ is a right Hilbert B-module $\mathcal{E}_B$ together with a left action of A on $\mathcal{E}$ given by a $*$-homomorphism $\phi_A : A\to\mathcal{L}(\mathcal{E})$, $a\cdot\xi = \phi_A(a)\xi$ for $a\in A$ and $\xi\in\mathcal{E}$. We may occasionally write, ${}_A\mathcal{E}_B$ to denote an $A-B$ C^*-correspondence and ϕ instead of ϕ_A. Furthermore, if ${}_{A_1}\mathcal{E}_{B_1}$ and ${}_{A_2}\mathcal{F}_{B_2}$ are C^*-correspondences, then a *morphism* $(\pi_1,T,\pi_2):\mathcal{E}\to\mathcal{F}$ consists of $*$-homomorphisms $\pi_i : A_i\to B_i$ and a linear map $T:\mathcal{E}\to\mathcal{F}$ satisfying

(i) $\pi_2(\langle\xi,\eta\rangle_{A_2}) = \langle T(\xi),T(\eta)\rangle_{B_2}$
(ii) $T(\phi_{A_1}(a_1)\xi) = \phi_{B_1}(\pi_1(a_1))T(\xi)$
(iii) $T(\xi)\pi_2(a_2) = T(\xi a_2)$

for all $\xi,\eta\in\mathcal{E}$ and $a_i\in A_i$.

Notation 2.5. When $A = B$, we refer to ${}_A\mathcal{E}_A$ as a C^*-correspondence over A. For $\mathcal{E}$ a C^*-correspondence over A and $\mathcal{F}$ a C^*-correspondence over B, a morphism $(\pi,T,\pi):\mathcal{E}\to\mathcal{F}$ will be denoted by (T,π).

Definition 2.6. [56] If $\mathcal{F}$ is the Hilbert module ${}_CC_C$ where C is a C^*-algebra with the inner product $\langle x,y\rangle_B = x^*y$ then call a morphism $(T,\pi):{}_A\mathcal{E}_B\to C$ of Hilbert modules a *representation* of ${}_A\mathcal{E}_B$ into C.

Remark 2.7. Note that a representation of ${}_A\mathcal{E}_B$ need only satisfying (i) and (ii) of definition 2.4 as it was unnecessary to require (iii) since

$$\begin{aligned}||T(\xi)\pi(a) - T(\xi a)||^2 &= ||(T(\xi)\pi(a) - T(\xi a))^*(T(\xi)\pi(a) - T(\xi a))|| \\ &= ||\pi(a)^*T(\xi)^*T(\xi)\pi(a) - T(\xi a)^*T(\xi)\pi(a) \\ &\quad - \pi(a)^*T(\xi)^*T(\xi a) + T(\xi a)^*T(\xi a)|| \\ &= ||\pi(a^*\langle\xi,\xi\rangle a - \langle\xi a,\xi\rangle a - a^*\langle\xi,\xi a\rangle + \langle\xi a,\xi a\rangle)|| \\ &= ||\pi(a^*\langle\xi,\xi\rangle a - a^*\langle\xi,\xi\rangle a - a^*\langle\xi,\xi\rangle a + a^*\langle\xi,\xi\rangle a)|| = 0\end{aligned}$$

by condition (i).

A morphism of Hilbert modules $(T,\pi) : \mathcal{E} \to \mathcal{F}$ yields a $*$-homomorphism $\Psi_T : \mathcal{K}(\mathcal{E}) \to \mathcal{K}(\mathcal{F})$ by

$$\Psi_T(\theta^{\mathcal{E}}_{\xi,\eta}) = \theta^{\mathcal{F}}_{T(\xi),T(\eta)}$$

for $\xi,\eta \in \mathcal{E}$ and if $(S,\sigma) : \mathcal{D} \to \mathcal{E}$, and $(T,\pi) : \mathcal{E} \to \mathcal{F}$ are morphisms of Hilbert modules then $\Psi_T \circ \Psi_S = \Psi_{T\circ S}$. In the case where $\mathcal{F} = B$ a C^*-algebra, we may first identify $\mathcal{K}(B)$ as B, and a representation (T,π) of $\mathcal{E}$ in a C^*-algebra B yields a $*$-homomorphism $\Psi_T : \mathcal{K}(\mathcal{E}) \to B$ given by

$$\Psi_T(\theta_{\xi,\eta}) = T(\xi)T(\eta)^*.$$

Definition 2.8. [56] For a C^*-correspondence $\mathcal{E}$ over A, denote the ideal $\phi^{-1}(\mathcal{K}(\mathcal{E}))$ of A by $J(\mathcal{E})$, and let $J_{\mathcal{E}} = J(\mathcal{E}) \cap (\ker\phi)^\perp$ where $(\ker\phi)^\perp$ is the ideal $\{a \in A \,\big|\, ab = 0 \text{ for all } b \in \ker\phi\}$. If ${}_A\mathcal{E}_A$ and ${}_B\mathcal{F}_B$ are C^*-correspondences over A and B respectively and $K \lhd J(\mathcal{E})$, a morphism $(T,\pi) : \mathcal{E} \to \mathcal{F}$ is called *coisometric on* K if

$$\Psi_T(\phi_A(a)) = \phi_B(\pi(a))$$

for all $a \in K$, or just *coisometric*, if $K = J(\mathcal{E})$.

Notation 2.9. We denote $C^*(T,\pi)$ to be the C^*-algebra generated by $T(\mathcal{E})$ and $\pi(A)$ where $(T,\pi) : \mathcal{E} \to B$ is a representation of ${}_A\mathcal{E}_A$ in a C^*-algebra B. Furthermore, if $\rho : B \to C$ is a $*$-homomorphism of C^*-algebras, then $\rho \circ (T,\pi)$ denotes the representation $(\rho \circ T, \rho \circ \pi)$ of $\mathcal{E}$.

Definition 2.10. [56] A morphism $(T_{\mathcal{E}},\pi_{\mathcal{E}})$ coisometric on an ideal K is said to be *universal* if whenever $(T,\pi) : \mathcal{E} \to B$ is a representation coisometric on K, there exists a $*$-homomorphism $\rho : C^*(T_{\mathcal{E}},\pi_{\mathcal{E}}) \to B$ with $(T,\pi) = \rho \circ (T_{\mathcal{E}},\pi_{\mathcal{E}})$. The universal C^*-algebra $C^*(T_{\mathcal{E}},\pi_{\mathcal{E}})$ is called the *relative Cuntz-Pimsner algebra* of $\mathcal{E}$ determined by the ideal K and denoted by $\mathcal{O}(K,\mathcal{E})$. If $K = 0$, then $\mathcal{O}(K,\mathcal{E})$ is denoted by $\mathcal{T}(\mathcal{E})$ and called the *universal Toeplitz C^*-algebra* for $\mathcal{E}$. We denote $\mathcal{O}(J_{\mathcal{E}},\mathcal{E})$ by $\mathcal{O}_{\mathcal{E}}$.

In [38], Katsura provides a sufficient condition to guarantee the nuclearity of the Cuntz-Pimsner algebra $\mathcal{O}_{\mathcal{E}}$:

Proposition 2.11. [38, Corollary 7.4] Let $\mathcal{E}$ be a C^*-correspondence over C^*-algebra A. If A is nuclear, then $\mathcal{O}_{\mathcal{E}}$ is nuclear.

2.2. **Topological Quivers.** Many C^*-algebras are achieved as graph C^*-algebras [43, 51] such as 0, $\mathbb{C}^d$ ($d \in \mathbb{N}$), $M_n(\mathbb{C})$ ($n \in \mathbb{N}$), $\mathcal{K}$ (the compact operators on a separable Hilbert space), the Cuntz algebras $\mathcal{O}_n$ ($n \in \mathbb{N}$) and the Toeplitz Algebra $\mathcal{T}$. It was proven in [25] for sinkless graph G and generalized in [8] that $C^*(G)$ is a (relative) Cuntz-Pimsner algebra.

A certain notion of "continuous graph C^*-algebra" exists when X and E are infinite (perhaps nondiscrete) sets, yet a few extra manageable properties for X, and E are needed. This notion is what is called a topological quiver which may be found in [9, 10, 11, 56].

Definition 2.12. [56] A *topological quiver* (or *topological directed graph*) $Q = (X, E, Y, r, s, \lambda)$ is a diagram

$$\begin{tikzcd} X & E \arrow[l, "s"'] \arrow[r, "r"] & Y \end{tikzcd}$$

where X, E, and Y are second countable locally compact Hausdorff spaces, r and s are continuous maps with r open, along with a family $\lambda = \{\lambda_y | y \in Y\}$ of Radon measures on E satisfying

(1) $\operatorname{supp} \lambda_y = r^{-1}(y)$ for all $y \in Y$, and
(2) $y \mapsto \lambda_y(f) = \int_E f(\alpha) d\lambda_y(\alpha) \in C_c(Y)$ for $f \in C_c(E)$.

If $X = Y$ then write $Q = (X, E, r, s, \lambda)$ in lieu of (X, E, X, r, s, λ).

Remark 2.13. It is important to remark that in the literature pertaining to graph C^*-algebras and their generalizations, various authors interchange the roles of the maps r and s in their definitions. Definition 2.12 agrees with that used in most papers on graph C^*-algebras (cf. [2, 3, 7, 9, 10, 46, 47, 56]). However, Definition 2.12 differs from that used in the higher rank graph algebras of Kumjian and Pask [44, 45] and from that used in the topological graph algebras of Katsura [39, 40, 41, 42] where Katsura uses the term topological graph to denote a more restrictive concept.

Given a topological quiver $Q = (X, E, Y, r, s, \lambda)$, one may associate a correspondence $\mathcal{E}_Q$ of the C^*-algebra $C_0(X)$ to the C^*-algebra $C_0(Y)$. Define left and right actions

$$(a \cdot \xi \cdot b)(e) = a(s(e))\xi(e)b(r(e))$$

by $C_0(X)$ and $C_0(Y)$ respectively on $C_c(E)$. Furthermore, define the $C_c(Y)$-valued inner product

$$\langle \xi, \eta \rangle(y) = \int_{r^{-1}(y)} \overline{\xi(\alpha)} \eta(\alpha) d\lambda_y(\alpha)$$

for $\xi, \eta \in C_c(E)$, $y \in Y$, and let $\mathcal{E}_Q$ be the completion of $C_c(E)$ with respect to the norm

$$||\xi|| = ||\langle \xi, \xi \rangle^{1/2}||_\infty = ||\lambda_y(|\xi|^2)||_\infty^{1/2}.$$

Definition 2.14. Given topological quiver Q over a space X, define the C^*-algebra, $C^*(Q)$ associated with Q to be the Cuntz-Pimnser C^*-algebra $\mathcal{O}_{\mathcal{E}_Q}$ of the correspondence $\mathcal{E}_Q$ over $A = C_0(X)$.

Proposition 2.15. [53, Proposition 2.21] If the left action, ϕ, is injective, then the universal map $i_A : A \to \mathcal{O}_{\mathcal{E}_Q}$ is injective.

Proposition 2.16. $C^*(Q)$ is nuclear for any topological quiver Q.
Proof. Since abelian C^*-algebras are nuclear, so is $A = C_0(X)$ and hence by Proposition 2.11, $C^*(Q)$ is nuclear.

□

Example 2.17.

(1) Let $G = (X, E, r, s)$ be a directed graph and let λ_y be counting measure on $r^{-1}(y)$. Then G along with this family of Radon measures λ on E becomes a topological quiver $Q = (X, E, r, s, \lambda)$. As we have already mentioned, $C^*(G) \cong C^*(Q)$.
(2) In [39], Katsura uses topological graph $G = (X, E, r, d)$ to be the more restrictive situation using two continuous maps $r, d : E \to X$ with d a local homeomorphism. We may set λ_x to be counting measure on $d^{-1}(x)$ since $d^{-1}(x)$ must be a discrete space. Then $Q = (X, E, d, r, \lambda)$ is a topological quiver. Furthermore, Katsura defines an associated C^*-algebra $C^*(G)$ which is isomorphic to $C^*(Q)$ [39, Definition 2.9 and Definition 2.10].
(3) In [7], Brenken defines a C^*-algebra associated to a closed relation, $\alpha \subset X \times X$, on a topological space X with π_i the usual projection map onto the i-th coordinate. There is a Radon measure μ_x such that supp $\mu_x = \pi_2^{-1}(x)$ for all $x \in X$. The Cuntz-Pimsner algebra of the associated correspondence, $C^*(\alpha)$ is isomorphic to $C^*(Q)$ where Q is the topological quiver $(X, \alpha, \pi_2, \pi_1, \mu)$. Such topological quivers are referred to as *topological relations.*

Theorem 2.18. Let $Q = (X, E, r, s, \lambda)$ be a topological quiver with X compact. If $\mathcal{E}_Q$ has a finite orthonormal basis $\{u_i\}_{i=1}^n \subset C_c(E)$ and ϕ is injective then $C^*(Q)$ is the universal C^*-algebra generated by $C(X)$ and elements $\{S_\xi\}_{\xi \in \mathfrak{I}}$, where $\mathfrak{I}$ is the module generated by $\{u_i\}_{i=1}^n$, satisfying the relations:

(1) $S_{\alpha\xi+\beta\eta} = \alpha S_\xi + \beta S_\eta$,
(2) $a \cdot S_\xi \cdot b = S_{a\cdot\xi\cdot b}$,
(3) $S_{u_i}^* S_{u_j} = \delta_i^j 1 \in C(X)$, and
(4) $\sum_{i=1}^n S_{u_i} S_{u_i}^* = 1$

for all $\alpha, \beta \in \mathbb{C}$, $a, b \in C(X)$ and $\xi, \eta \in \mathfrak{I}$. Moreover, the Toeplitz-Pimsner algebra $\mathcal{T}(Q) := \mathcal{T}(\mathcal{E}_Q)$ is the universal C^*-algebra satisfying (1)-(3).
Proof. Let $\mathcal{A}$ be the universal C^*-algebra generated by $C(X)$ and $\{S_\xi \,|\, \xi \in \mathfrak{I}\}$ satisfying the relations:

(1) $S_{\alpha\xi+\beta\eta} = \alpha S_\xi + \beta S_\eta$,
(2) $a \cdot S_\xi \cdot b = S_{a\cdot\xi\cdot b}$,
(3) $S_{u_i}^* S_{u_j} = \delta_i^j 1$, and
(4) $\sum_{i=1}^n S_{u_i} S_{u_i}^* = 1$

for all $\alpha, \beta \in \mathbb{C}$, $a, b \in C(X)$ and $\xi, \eta \in \mathfrak{I}$. To show the existence of $\mathcal{A}$, calculate

$$\begin{aligned} ||S_\xi|| &= ||S_{\sum_{i=1}^n u_i \langle u_i, \xi \rangle}|| \\ &= ||\sum_{i=1}^n S_{u_i} \langle u_i, \xi \rangle|| \\ &\leq \sum_{i=1}^n ||S_{u_i}|| ||\langle u_i, \xi \rangle||_\infty \end{aligned}$$

for each $\xi \in \mathfrak{I}$ and furthermore, $||S_{u_i}||^2 = ||S_{u_i}^* S_{u_i}|| = 1$. Since $||\langle u_i, \xi \rangle||_\infty$ is bounded in $C(X)$, $\mathcal{A}$ exists as a universal C^*-algebra.

Now if $a \in C(X)$ then

$$\sum_{i=1}^n \theta_{a \cdot u_i, u_i}(\xi)(e) = \sum_{i=1}^n (a \cdot u_i) \langle u_i, \xi \rangle (e) = a(s(e))\xi(e) = (\phi(a)\xi)(e);$$

that is, $J(\mathcal{E}_Q) = C(X)$. Then let (T_u, π_u) be the universal representation of $\mathcal{E}_Q$ in $C^*(Q)$ coisometric on $J_{\mathcal{E}_Q} = J(\mathcal{E}_Q)$ (since ϕ is injective.) Setting $S_\xi = T_u(\xi)$, we have $S_{\alpha\xi + \beta\eta} = \alpha S_\xi + \beta S_\eta$, $a \cdot S_\xi \cdot b = S_{a \cdot \xi \cdot b}$, and $S_{u_i}^* S_{u_j} = \langle u_i, u_j \rangle = \delta_i^j 1$ for all $\alpha, \beta \in \mathbb{C}$, $a, b \in C(X)$ and $\xi, \eta \in \mathfrak{I}$. Finally, the reconstruction formula can easily be rewritten as

$$\sum_{i=1}^n \theta_{u_i, u_i} = 1 \in C(X) = J_{\mathcal{E}_Q};$$

hence, (4) is evident. Thus, there exists a homomorphism $\rho : \mathcal{A} \to C^*(Q)$ such that

$$S_\xi \mapsto T_u(\xi)$$

and

$$a \mapsto \pi_u(a)$$

for each $\xi \in \mathfrak{I}$ and $a \in C(X)$.

Conversely, given $\{S_\xi\}_{\xi \in \mathfrak{I}}$ satisfying (1)-(4), we construct a coisometric representation (T, π) of $\mathcal{E}_Q$ into $\mathcal{A}$; that is, let $T(\xi) = S_\xi$ for each $\xi \in \mathcal{E}_Q (= \mathfrak{I}$, since $\{u_i\}_{i=1}^n$ is a basis) and let π be the inclusion of $C(X)$ into the universal C^*-algebra stated above. With this in mind, (1) and (2) imply that T is a linear map that satisfies

$$T(a \cdot \xi \cdot b) = a \cdot T(\xi) \cdot b.$$

Furthermore, if $\xi, \eta \in \mathcal{E}_Q$ then

$$\begin{aligned} T(\xi)^*T(\eta) &= T(\sum_{i=1}^{n} u_i\langle u_i, \xi\rangle)^* T(\sum_{j=1}^{n} u_j\langle u_j, \eta\rangle) \\ &= [\sum_{i=1}^{n} \langle \xi, u_i\rangle S_{u_i}^*][\sum_{j=1}^{n} S_{u_j}\langle u_j, \eta\rangle] \\ &= \sum_{i=1}^{n} \langle \xi, u_i\rangle\langle u_i, \eta\rangle = \sum_{i=1}^{n} \langle u_i\langle u_i, \xi\rangle, \eta\rangle \\ &= \langle \sum_{i=1}^{n} u_i\langle u_i, \xi\rangle, \eta\rangle = \langle \xi, \eta\rangle. \end{aligned}$$

Finally, if $a \in C(X)$ then

$$\Psi_T(\phi(a)) = \Psi_T(\sum_{i=1}^{n} \theta_{a\cdot u_i, u_i}) = \sum_{i=1}^{n} S_{a\cdot u_i} S_{u_i}^* = a\sum_{i=1}^{n} S_i S_i^* = a.$$

Thus, $\Psi_T(\phi(a)) = a$ and hence, (T, π) is coisometric. Thus, there exist a homomorphism $\tau : C^*(Q) \to \mathcal{A}$ such that $\tau \circ (T_u, \pi_u) \to (T, \pi)$ where (T_u, π_u) is the universal coisometric representation of $\mathcal{E}_Q$ in $C^*(Q)$. Thus,

$$\tau \circ \rho(S_\xi) = \tau(T_u(\xi)) = T(\xi) = S_\xi$$

and

$$\tau \circ \rho(a) = \tau(\pi_u(a)) = \pi(a) = a$$

for each $\xi \in \mathcal{E}_Q$ and $a \in A$; that is, ρ and τ are isomorphisms.

□

Remark 2.19. The existence of a basis $\{u_i\}_{i=1}^n$ forces $\sum_{i=1}^n \theta_{u_i,u_i} = 1$ and hence, X must be compact and ϕ must be injective.

2.3. **Topological Quiver Isomorphisms.** A notion for a topological quiver isomorphism based on the definition of a directed graph isomorphism is introduced.

Definition 2.20. Let $Q = (X, E, Y, r, s, \lambda)$ and $Q' = (X', E', Y', r', s', \lambda')$ be topological quivers. If there exist homeomorphisms $\varphi_1 : X \to X'$, $\varphi_2 : Y \to Y'$, and $\psi : E \to E'$ where ψ is measurable and there exists a family of constants $\{h_y > 0 \,|\, y \in Y\}$ such that

1. $\varphi_1(s(e)) = s'(\psi(e))$
2. $\varphi_2(r(e)) = r'(\psi(e))$
3. $\lambda'_{\varphi(y)}(B) = \int_B h_y \, d(\lambda_y \circ \psi^{-1})$

for each $e \in E$, measurable $B \subseteq E$ and $y \in Y$, then say Q and Q' are *isomorphic* and write $Q \cong Q'$.

Remark 2.21. Recall that a measurable map $\psi : E \to E'$ is one that satisfies $\psi^{-1}(B')$ is measurable in E for each measurable subset $B' \subseteq E'$ (see [67].) For any homeomorphism $\psi : E \to E'$ and compact $K \subset E'$, assume $\psi^{-1}(K)$ is covered by the open collection $\{U_i\}_{i\in\mathcal{I}}$. Then $\psi^{-1}(K) \subset \cup_{i\in\mathcal{I}} U_i$ implies $K \subset \cup_{i\in\mathcal{I}}\psi(U_i)$. Hence, K is covered by the open sets $\psi(U_i)$ and so, $K \subset \cup_{j=1}^{n}\psi(U_{i_j})$ for some open subcover $\{\psi(U_{i_j})\}_{j=1}^{n}$. Finally, $\psi^{-1}(K) \subset \cup_{j=1}^{n} U_{i_j}$; that is, $\psi^{-1}(K)$ is compact and thus, ψ is a proper map. Note assuming ψ is open, injective and continuous is enough to show that ψ is a proper map.

Theorem 2.22. Let $Q = (X, E, r, s, \lambda)$ and $Q' = (X', E', r', s', \lambda')$ be topological quivers. If $Q \cong Q'$ by (φ, ψ, h), then $\mathcal{O}(K, \mathcal{E}_Q) \cong \mathcal{O}(K', \mathcal{E}_{Q'})$ for any closed ideal K' of $J(\mathcal{E}_Q)$ where $K = \varphi^{\#}(K')$; that is, given a homeomorphism $\varphi : X \to X'$, a measurable homeomorphism $\psi : E \to E'$ and a family of constants $\{h_y > 0 \,|\, y \in X\}$ satisfying

1. $\varphi_1(s(e)) = s'(\psi(e))$
2. $\varphi_2(r(e)) = r'(\psi(e))$
3. $\lambda'_{\varphi(y)}(B) = \int_B h_y \, d(\lambda_y \circ \psi^{-1})$

for each $e \in E$, measurable $B \subseteq E$ and $y \in Y$, then $\mathcal{O}(K, \mathcal{E}_Q) \cong \mathcal{O}(K', \mathcal{E}_{Q'})$.

Proof. Define $T_0 : C_c(E') \to C_c(E)$ and $\pi : C_0(X') \to C_0(X)$ by

$$T_0(\xi')(e) = h_{r(e)}^{1/2}\xi'(\psi(e)) \qquad \text{and} \qquad \pi(a') = a' \circ \varphi$$

for all $\xi' \in C_c(E')$ and $a' \in C_0(X')$. Since ψ is proper, $T_0(\xi') \in C_c(E)$ for each $\xi' \in C_c(E')$. For $\xi', \eta' \in C_c(E')$, $\alpha, \beta \in \mathbb{C}$ and $e \in E$,

$$\begin{aligned} T_0(\alpha\xi' + \beta\eta')(e) &= h_{r(e)}^{1/2}(\alpha\xi' + \beta\eta')(\psi(e)) \\ &= h_{r(e)}^{1/2}(\alpha\xi'(\psi(e)) + \beta\eta'(\psi(e))) \\ &= \alpha h_{r(e)}^{1/2}\xi'(\psi(e)) + \beta h_{r(e)}^{1/2}\eta'(\psi(e)) \\ &= (\alpha T_0(\xi') + \beta T_0(\eta'))(e); \end{aligned}$$

that is, T_0 is a linear map, and hence there exists a lift to a linear map $T : \mathcal{E}_{Q'} \to \mathcal{E}_Q$ such that $T\big|_{C_c(E')} = T_0$. We now claim that (T, π) is a coisometric morphism of correspondence $\mathcal{E}_{Q'}$ to correspondence $\mathcal{E}_Q$.

To see this claim, let $y \in X$, $e \in E$, $\xi', \eta' \in C_c(E')$ and $a' \in C_0(X')$:

$$
\begin{aligned}
(1)\ \pi(\langle \xi', \eta' \rangle)(y) &= \langle \xi', \eta' \rangle(\varphi(y)) = \int \overline{\xi'(e')}\eta'(e')\, d\lambda'_{\varphi(y)}(e') \\
&= \int \overline{(\xi' \circ \psi)(\psi^{-1}(e'))}(\eta' \circ \psi)(\psi^{-1}(e'))h_{r(e)}\, d(\lambda_y \circ \psi^{-1})(e') \\
&= \int \overline{h^{1/2}_{r(e)}(\xi' \circ \psi)(e_0)}h^{1/2}_{r(e)}(\eta' \circ \psi)(e_0)\, d\lambda_y(e_0) \\
&= \langle T(\xi'), T(\eta') \rangle(y)
\end{aligned}
$$

$$
\begin{aligned}
(2)\ T(\xi' \cdot a')(e) &= h^{1/2}_{r(e)}(\xi' \cdot a')(\psi(e)) \\
&= h^{1/2}_{r(e)}(\xi' \circ \psi)(e)(a'(r'(\psi(e)))) \\
&= T(\xi')(e)(a' \circ \varphi)(r(e)) \\
&= T(\xi')\pi(a')(e)
\end{aligned}
$$

and similarly,

$$(3)\ T(\phi(a')\xi')(e) = \pi(a')T(\xi')(e).$$

Finally, (4) Suppose $a' \in J(\mathcal{E}_{Q'}) = (\phi')^{-1}(\mathcal{K}(\mathcal{E}_{Q'}))$. Then for every $\epsilon > 0$, there exist θ_ϵ of the form

$$\theta_\epsilon = \sum_{k=1}^{n_\epsilon} \theta_{\xi'_k, \eta'_k} \in \mathcal{K}(\mathcal{E}_{Q'})$$

where $\xi'_k, \eta'_k \in C_c(E')$ such that $||\theta_\epsilon - \phi'(a')|| < \epsilon$. To show that (T, π) is coisometric, it is enough to show that $\Psi_T(\phi'(a')) = \phi(\pi(a'))$ for all $a' = \theta_{\xi',\eta'}$ where $\xi', \eta' \in C_c(E')$. To this end, let $\zeta \in C_c(E)$ and $e \in E$. Then

$$
\begin{aligned}
\phi(\pi(\theta_{\xi',\eta'}))\zeta(e) &= (\theta_{\xi',\eta'} \circ \varphi)(s(e))\zeta(e) = \theta_{\xi',\eta'}(s(\psi(e)))\zeta(e) \\
&= (\theta_{\xi',\eta'} \cdot (\zeta \circ \psi^{-1}))(\psi(e)) = \xi'\langle \eta', \zeta \circ \psi^{-1} \rangle(\psi(e)) \\
&= \xi'(\psi(e)) \int \overline{\eta'(e')}(\zeta \circ \psi^{-1})(e')\, d\lambda'_{r'(\psi(e))}(e') \\
&= \xi'(\psi(e)) \int \overline{(\eta' \circ \psi)(\psi^{-1}(e'))}(\zeta \circ \psi^{-1})(e')\, d\lambda'_{\varphi(r(e))}(e') \\
&= \xi'(\psi(e)) \int \overline{(\eta' \circ \psi)(\psi^{-1}(e'))}(\zeta \circ \psi^{-1})(e')\, d\lambda_{r(e)}(\psi^{-1}(e')) \\
&= \xi'(\psi(e)) \int \overline{(\eta' \circ \psi)(e_0)}\zeta(e_0)h_{r(e)}\, d\lambda_{r(e)}(e_0)) \\
&= h^{1/2}_{r(e)}(\xi' \circ \psi)(e)\langle h^{1/2}_{r(e)}(\eta' \circ \psi), \zeta \rangle(r(e)) \\
&= T(\xi') \cdot \langle T(\eta'), \zeta \rangle(e) \\
&= \phi'(\theta_{T(\xi'),T(\eta')})\zeta(e) \\
&= \Psi_T(\phi'(\theta_{\xi',\eta'}))\zeta(e).
\end{aligned}
$$

Hence, (T,π) is coisometric.

Now take the universal representations (T_K,π_K) and $(T_{K'},\pi_{K'})$ that are coisometric on $K=\varphi^{\#}(K')=\pi(K')$ and K', respectively. First, we need to show that K is a closed ideal of $J(\mathcal{E}_Q)$. Note it is easy to see that $\pi(a)\in\phi^{-1}(\mathcal{K}(\mathcal{E}_Q))$ for each $a\in J(\mathcal{E}_{Q'})$ from the long calculation above and so, $\pi(J(\mathcal{E}_{Q'}))\subseteq J(\mathcal{E}_Q)$. Furthermore, since $C_c(E)$ is in the image of T, we must have $J(\mathcal{E}_Q)=\pi(J(\mathcal{E}_{Q'}))$. If $a\in J(\mathcal{E}_Q)$ and $b\in K$, then there exist $a'\in J(\mathcal{E}_{Q'})$ and $b'\in K'$ such that $a=\pi(a')$ and $b=\pi(b')$. So

$$ab=\pi(a'b')\in\pi(K')=K.$$

Likewise for $ba\in K$ and closure follows from the fact that φ is a homeomorphism.

We now claim that $(T_K,\pi_K)\circ(T,\pi)$ is a morphism that is coisometric on K'. To see this, let ϕ_K be the left action homomorphism and $a'\in K'$, then $\pi(a')=\varphi^{\#}(a')\in K$ and

$$\Psi_{T_K\circ T}(\phi'(a'))=\Psi_{T_K}(\phi(\pi(a')))=\phi_K(\pi_K\circ\pi(a')).$$

Now by the universal property of $(T_{K'},\pi_{K'})$, there exists a $*$-homomorphism $\rho:\mathcal{O}(K',\mathcal{E}_{Q'})\to\mathcal{O}(K,\mathcal{E}_Q)$ satisfying

$$(T_K,\pi_K)\circ(T,\pi)=\rho\circ(T_{K'},\pi_{K'}).$$

We may do precisely the same with $(T^{-1},\pi^{-1}):\mathcal{E}_Q\to\mathcal{E}_{Q'}$ defined by $T_0^{-1}(\xi)(e')=h_{r'(e')}^{-1/2}(\xi\circ\psi^{-1})(e')$ for each $e'\in E'$ and $\xi\in C_c(E)$ with the lift $T^{-1}:\mathcal{E}_Q\to\mathcal{E}_{Q'}$ satisfying $T^{-1}\Big|_{C_c(E)}=T_0^{-1}$. Also, $\pi^{-1}(a)=a\circ\varphi^{-1}$ for each $a\in C_0(X)$. Furthermore, note $K'=(\varphi^{-1})^{\#}(K)$. Thus, by the universal property of (T_K,π_K), there exists a $*$-homomorphism $\tau:\mathcal{O}(K,\mathcal{E}_Q)\to\mathcal{O}(K',\mathcal{E}_{Q'})$ satisfying

$$(T_{K'},\pi_{K'})\circ(T^{-1},\pi^{-1})=\tau\circ(T_K,\pi_K)$$

and we note $\rho\circ\tau=id_{\mathcal{O}(K,\mathcal{E}_Q)}$ and $\tau\circ\rho=id_{\mathcal{O}(K',\mathcal{E}_{Q'})}$. Hence, $\mathcal{O}(K,\mathcal{E}_Q)\cong\mathcal{O}(K',\mathcal{E}_{Q'})$. □

Corollary 2.23. With the notation of Theorem 2.22, if $Q\cong Q'$ then $\mathcal{T}(Q)\cong\mathcal{T}(Q')$ and $C^*(Q)\cong C^*(Q')$.

Proof. We need only show $\pi(J_{\mathcal{E}_{Q'}})=J_{\mathcal{E}_Q}$. So given any $b\in\ker\phi$, $\xi'\in\mathcal{E}_{Q'}$ and $e'\in E'$,

$$\begin{aligned}\phi'(b\circ\varphi^{-1})\xi'(e')&=b(\varphi^{-1}(s'(e')))\xi'(e')\\&=b(s(\psi^{-1}(e')))(\xi'\circ\psi)(\psi^{-1}(e'))\\&=b\cdot(\xi\circ\psi)(\psi^{-1}(e'))=0\end{aligned}$$

since $\phi(b)=0$, and so $b\circ\varphi^{-1}\in\ker\phi'$. Furthermore, for each $a'\in J_{\mathcal{E}_{Q'}}$,

$$\phi(\pi(a'))b=(a'\circ\varphi)b=(a(b\circ\varphi^{-1}))\circ\varphi=0.$$

Hence, $\pi(a)\in(\ker\phi')^{\perp}$ and recall from the proof of the previous theorem that $\pi(J(\mathcal{E}_{Q'}))=J(\mathcal{E}_Q)$, so $\pi(J_{\mathcal{E}_{Q'}})\subseteq J_{\mathcal{E}_Q}$. Conversely, let $a\in J_{\mathcal{E}_Q}$. Then there exists $a'\in J(\mathcal{E}_{Q'})$ such that $\pi(a')=a$, and given any $b'\in\ker\phi'$,

$$\pi(a'b')=a\pi(b')=0$$

since $\pi(b') \in \ker\phi$. To see this, let $\xi \in \mathcal{E}_Q$ and $e \in E$. Then

$$\begin{aligned}\phi(\pi(b'))\xi(e) &= b'(\varphi(s(e)))\xi(e)\\ &= b'(s'(\psi(e)))(\xi\circ\psi^{-1})(\psi(e))\\ &= \phi'(b')(\xi\circ\psi^{-1})(\psi(e)) = 0 \qquad (\text{since } b' \in \ker\phi'.)\end{aligned}$$

Hence, $a'b' = 0$ and thus, $a' \in (\ker\phi')^{\perp}$, as desired.

□

3. Topological Group Quivers

3.1. **Definitions.** We now investigate a particular class of topological quivers related to a locally compact group and then form and study its associated C^*-algebra. This construction leads us to many interesting results including a presentation of the C^*-algebra, and the spatial structure of the C^*-algebra.

Definition 3.1. Let Γ be a (second countable) locally compact group and let $\alpha, \beta \in \mathrm{End}(\Gamma)$ be continuous. Define the closed subgroup, $\Omega_{\alpha,\beta}(\Gamma)$, of $\Gamma\times\Gamma$,

$$\Omega_{\alpha,\beta}(\Gamma) = \{(x,y) \in \Gamma\times\Gamma \,\big|\, \alpha(y) = \beta(x)\}$$

and let $Q_{\alpha,\beta}(\Gamma) = (\Gamma, \Omega_{\alpha,\beta}(\Gamma), r, s, \lambda)$ where r and s are the group homomorphisms defined by

$$r(x,y) = x \qquad \text{and} \qquad s(x,y) = y$$

for each $(x,y) \in \Omega_{\alpha,\beta}(\Gamma)$ and λ_x for $x \in \Gamma$ is the measure on

$$r^{-1}(x) = \{x\}\times\alpha^{-1}(\beta(x))$$

defined by

$$\lambda_x(B) = \mu(y^{-1}s(B\cap r^{-1}(x))\cap\ker\alpha) \qquad (\text{for any } y \in \alpha^{-1}(\beta(x)))$$

for each measurable $B \subseteq \Omega_{\alpha,\beta}(\Gamma)$ where μ is a left Haar measure (normalized if possible) on $r^{-1}(1_\Gamma) = \{1\}\times\ker\alpha$ (a closed normal subgroup of $\Gamma\times\Gamma$; hence, a locally compact group). Note if $r^{-1}(x) = \emptyset$ then $\alpha^{-1}(\beta(x)) = \emptyset$ and so $\lambda_x = 0$. This measure is well-defined. If $y_1, y_2 \in \alpha^{-1}(\beta(x))$ then $y_2^{-1}y_1 \in \ker\alpha$ and $y_2^{-1}y_1\ker\alpha = \ker\alpha$. Hence,

$$\begin{aligned}\mu(y_1^{-1}s(B\cap r^{-1}(x))\cap\ker\alpha) &= \mu(y_2^{-1}y_1[y_1^{-1}s(B\cap r^{-1}(x))\cap\ker\alpha])\\ &= \mu([y_2^{-1}y_1y_1^{-1}s(B\cap r^{-1}(x))]\cap[y_2^{-1}y_1\ker\alpha])\\ &= \mu(y_2^{-1}s(B\cap r^{-1}(x))\cap\ker\alpha)\end{aligned}$$

for each measurable $B \subseteq \Omega_{\alpha,\beta}(\Gamma)$.

Furthermore, let $x \in \Gamma$. If $y \in \alpha^{-1}(\beta(x)) \neq \emptyset$ then for any $y_0 \in \alpha^{-1}(\beta(x))$,

$$\alpha(y^{-1}y_0) = \alpha(y)^{-1}\alpha(y_0) = \beta(x)^{-1}\beta(x) = 1_\Gamma;$$

hence, $y^{-1}y_0 \in \ker\alpha$ and $y_0 = yy^{-1}y_0 \in y\ker\alpha$. So $\alpha^{-1}(\beta(x)) \subseteq y\ker\alpha$ and if $y_0 = yz$ for some $z \in \ker\alpha$, then $\alpha(y_0) = \alpha(yz) = \alpha(y) = \beta(x)$. Thus, $\alpha^{-1}(\beta(x)) = y\ker\alpha$. So if $r^{-1}(x) = \emptyset$ then $\operatorname{supp}\lambda_x = \emptyset = r^{-1}(x)$ and if $(x,y) \in r^{-1}(x)$ then

$$\operatorname{supp}\lambda_x = \{x\}\times y\ker\alpha = \{x\}\times\alpha^{-1}(\beta(x)) = r^{-1}(x).$$

Moreover, the map

$$x \mapsto \int_{r^{-1}(x)} f(e)\, d\lambda_x(e) = \int_{\ker\alpha} f(x, yz)\, d\mu(z)$$

has (closed) support contained in $r(\operatorname{supp} f)$ (which is compact for each $f \in C_c(\Omega_{\alpha,\beta}(\Gamma))$.) Finally, the combined facts that α and β are continuous endomorphisms (hence, open maps) and f has compact support guarantees that this map is continuous.

Call $Q_{\alpha,\beta}(\Gamma)$ a *topological group relation.* Define $\mathcal{E}_{\alpha,\beta}(\Gamma)$ to be the $C_0(\Gamma)$-correspondence $\mathcal{E}_{Q_{\alpha,\beta}(\Gamma)}$ and form the Cuntz-Pimsner algebra

$$\mathcal{O}_{\alpha,\beta}(\Gamma) := C^*(Q_{\alpha,\beta}(\Gamma)) = \mathcal{O}(J_{\mathcal{E}_{\alpha,\beta}(\Gamma)}, \mathcal{E}_{\alpha,\beta}(\Gamma))$$

and the Toeplitz-Pimsner algebra

$$\mathcal{T}_{\alpha,\beta}(\Gamma) := \mathcal{T}(Q_{\alpha,\beta}(\Gamma)).$$

Remark 3.2. It will be implicitly assumed that Γ is second countable. Furthermore, since Γ is locally compact Hausdorff, $r^{-1}(x)$ is closed and locally compact. Moreover, whenever r is a local homeomorphism, $r^{-1}(x)$ is discrete and hence, λ_x is counting measure (normalized when $|\ker\alpha| < \infty$.)

Further note the nomenclature "topological group relation" is chosen because the topological quiver $Q_{\alpha,\beta}(\Gamma)$ is a topological relation as introduced in [7].

Remark 3.3. By Theorem 2.22, we see that the resulting C^*-algebra is independent of the left Haar measure chosen, since any two such left Haar measures will satisfy $\mu = c\mu'$ for some $c > 0$. Thus we may solely regard the normalized left Haar measure when available.

Lemma 3.4. If Γ is a compact group and α is a continuous endomorphism on Γ, then $\ker\alpha$ is either finite or an uncountable perfect compact set.

Proof. Assume $\ker\alpha = \{z_i\}_{i\in I}$ is not finite. Then since Γ is compact and $\ker\alpha$ is the preimage of a closed set, $\ker\alpha$ is compact. Furthermore, let $\lim z_n = z \in \ker\alpha$ where $\{z_n \neq z\}_{n\in\mathbb{N}}$ is a convergent subsequence. Then, for any $y \in \ker\alpha$,

$$y = \lim_n yz^{-1}z_n$$

but $y \neq yz^{-1}z_n$ for all n, so $\ker\alpha$ is a perfect set. The Baire Category Theorem implies any perfect compact space must be uncountable.

□

We will be mainly concerned with examples where $\ker\alpha$ is finite, but we shall give a non-trivial example when $|\ker\alpha| = \infty$ in a later subsection. If $|\ker\alpha| < \infty$ and $y \in \alpha^{-1}(\beta(x))$ then

$$|r^{-1}(x)| = |\{x\} \times \alpha^{-1}(\beta(x))| = |\alpha^{-1}(\beta(x))| = \{y\ker\alpha\} = |\ker\alpha| < \infty.$$

Hence, the inner product on $\mathcal{E}_{\alpha,\beta}(\Gamma)$ becomes

$$\langle \xi, \eta\rangle(x) = \frac{1}{|\ker\alpha|} \sum_{\alpha(y)=\beta(x)} \overline{\xi(x,y)}\eta(x,y).$$

Moreover, one may inquire,"When is $Q_{\alpha,\beta}(\Gamma)$ sourceless/sinkless?" This is, in fact, quite easy to see. We note $r(\Omega_{\alpha,\beta}(\Gamma)) = \alpha^{-1}(\beta(\Gamma))$ and $s(\Omega_{\alpha,\beta}(\Gamma)) = \beta^{-1}(\alpha(\Gamma))$. Hence, $Q_{\alpha,\beta}(\Gamma)$ is sourceless (or sinkless) if and only if $\alpha^{-1}(\beta(\Gamma)) = \Gamma$ (or $\beta^{-1}(\alpha(\Gamma)) = \Gamma$.) In particular, $Q_{\alpha,\beta}(\Gamma)$ is sinkless and sourceless if α and β are surjective.

Given a continuous function $\omega : \Gamma \to (0, \infty)$, define a family of Radon measures $\{\lambda_x^\omega \,|\, x \in \Gamma\}$ by

$$\lambda_x^\omega(B) = \int_B \omega(s(e))\, d\lambda_x(e)$$

for each measurable $B \subset \Omega_{\alpha,\beta}(\Gamma)$ and let

$$||\xi||_\omega^2 := ||\lambda_x^\omega(\xi)||_\infty = \sup_{x\in\Gamma}\{\int \omega(s(e))|\xi|^2(e)\, d\lambda_x(e)\}$$

for each $\xi \in \mathcal{E}_{\alpha,\beta}(\Gamma)$.

Lemma 3.5. Let Γ be a compact group. Then $||\cdot||_\omega$ is an equivalent norm of $||\cdot||$.
Proof. First, $||\cdot||_\omega$ is a seminorm and there exist $m, M > 0$ such that $0 < m \leq \omega(x) \leq M$ for each $x \in \Gamma$. Thus,

$$||\xi||_\omega^2 = \sup_{x\in\Gamma}\{\int \omega(s(e))|\xi|^2(e)\, d\lambda_x(e)\} \leq M \sup_{x\in\Gamma}\{\int |\xi|^2(e)\, d\lambda_x(e)\} = M||\xi||^2$$

for each $\xi \in \mathcal{E}_{\alpha,\beta}(\Gamma)$.

On the other hand, given $\xi \in \mathcal{E}_{\alpha,\beta}(\Gamma)$ there exists $x_0 \in \Gamma$ such that $||\xi||^2 = \langle \xi, \xi \rangle(x_0)$. So

$$\begin{aligned}
||\xi||_\omega^2 \geq \lambda_{x_0}^\omega(\xi) &= \int \omega(s(e))|\xi|^2(e)\, d\lambda_{x_0}(e) \\
&\geq m \int |\xi|^2(e)\, d\lambda_{x_0}(e) \\
&= m\langle \xi, \xi \rangle(x_0) \\
&= m||\xi||^2.
\end{aligned}$$

Thus, $m||\xi||^2 \leq ||\xi||_\omega^2 \leq M||\xi||^2$ for each $\xi \in \mathcal{E}_{\alpha,\beta}(\Gamma)$.

If $||\xi||_\omega = 0$ then $m||\xi||^2 \leq 0 \leq M||\xi||^2$. So $||\xi|| = 0$ and thus, $\xi = 0$. Hence, $||\cdot||_\omega$ is a norm.

□

Lemma 3.6. Suppose Γ is a compact group and $\alpha, \beta \in \operatorname{End}(\Gamma)$ are continuous such that α has finite kernel. Then the norm given by the inner product, $||\cdot||$, is equivalent to the sup-norm and $C(\Omega_{\alpha,\beta}(\Gamma))$ is complete with respect to $||\cdot||$. Moreover, $\mathcal{E}_{\alpha,\beta}(\Gamma) = C(\Omega_{\alpha,\beta}(\Gamma))$ is a full correspondence .
Proof. Note first that $\Omega_{\alpha,\beta}(\Gamma)$ is a closed subset of the compact set $\Gamma \times \Gamma$, hence $\Omega_{\alpha,\beta}(\Gamma)$ is compact. Note

$$|r^{-1}(x)| = \begin{cases} |\ker\alpha| & \text{if } r^{-1}(x) \neq \emptyset \\ 0 & \text{if } r^{-1}(x) = \emptyset \end{cases} < \infty$$

for each $x \in \Gamma$ and so for $\Gamma_0 = \{x \in \Gamma \,|\, r^{-1}(x) \neq \emptyset\}$,

$$\begin{aligned}
||\xi||^2 &= ||\langle \xi, \xi\rangle||_\infty \\
&= \max_{x\in\Gamma}\{\int_{r^{-1}(x)} |\xi|(e)\, d\lambda_x(e)\} \\
&= \max_{\Gamma_0}\{\frac{1}{|\ker\alpha|}\sum_{\alpha(y)=\beta(x)} |\xi|^2(x,y)\} \\
&= \frac{1}{|\ker\alpha|}\sum_{z\in\ker\alpha} \max_{x\in\Gamma_0}\{|\xi|^2(x,y_x z)\} \qquad \text{for any } y_x \in \alpha^{-1}(\beta(x)) \text{ (per } x \in \Gamma_0) \\
&\leq \frac{1}{|\ker\alpha|}\sum_{z\in\ker\alpha} \max_{e\in\Omega_{\alpha,\beta}(\Gamma)}\{|\xi|^2(e)\} \\
&= ||\xi||_\infty^2.
\end{aligned}$$

Next, $\frac{1}{|\ker\alpha|}\sum_{z\in\ker\alpha}|\xi|^2(x,yz) \leq ||\xi||^2$ for each $x \in \Gamma$ and $y \in \alpha^{-1}(\beta(x))$. Thus, $|\xi|^2(x,y) \leq |\ker\alpha| ||\xi||^2$ for each $(x,y) \in \Omega_{\alpha,\beta}(\Gamma)$ and so,

$$\frac{1}{|\ker\alpha|}||\xi||_\infty^2 \leq ||\xi||^2 \leq ||\xi||_\infty^2.$$

Since $C(\Omega_{\alpha,\beta}(\Gamma))$ is complete with respect to the sup-norm, we have that $C(\Omega_{\alpha,\beta}(\Gamma))$ is complete with respect to the norm given by the inner product.

Finally, note that $\mathcal{E}_{\alpha,\beta}(\Gamma)$ is full since given any $a \in C(\Gamma)$, it is possible to define $\xi, \eta \in C(\Omega_{\alpha,\beta}(\Gamma))$ by $\xi(x,y) = 1$ and $\eta(x,y) = a(x)$ for $(x,y) \in \Omega_{\alpha,\beta}(\Gamma)$ and note

$$\langle \xi, \eta\rangle(x) = \int \overline{\xi(e)}\eta(e)\, d\lambda_x(e) = a(x).$$

□

Remark 3.7. The assertion in the previous lemma concerning the full Hilbert module $\mathcal{E}_{\alpha,\beta}(\Gamma)$ remains true for infinite kernel. The proof is identical.

In many cases one can simplify α and β and yet preserve the structure of the topological quiver and hence, the resulting C^*-algebra.

Proposition 3.8. Let $\sigma \in \mathrm{Aut}(\Gamma)$ be continuous. Then

$$Q_{\alpha,\beta}(\Gamma) \cong Q_{\sigma\circ\alpha,\sigma\circ\beta}(\Gamma) \qquad \text{and} \qquad Q_{\alpha,\beta}(\Gamma) \cong Q_{\alpha\circ\sigma,\beta\circ\sigma}(\Gamma).$$

Thus,

$$\mathcal{O}_{\alpha,\beta}(\Gamma) \cong \mathcal{O}_{\sigma\circ\alpha,\sigma\circ\beta}(\Gamma), \qquad \mathcal{O}_{\alpha,\beta}(\Gamma) \cong \mathcal{O}_{\alpha\circ\sigma,\beta\circ\sigma}(\Gamma),$$

$$\mathcal{T}_{\alpha,\beta}(\Gamma) \cong \mathcal{T}_{\sigma\circ\alpha,\sigma\circ\beta}(\Gamma) \qquad \text{and} \qquad \mathcal{T}_{\alpha,\beta}(\Gamma) \cong \mathcal{T}_{\alpha\circ\sigma,\beta\circ\sigma}(\Gamma).$$

Proof. This is, in fact, a corollary of Theorem 2.22. For the first isomorphism, use $\varphi = id_\Gamma$ and $\psi = id_{\Omega_{\alpha,\beta}(\Gamma)}$ while noting

$$(x,y) \in \Omega_{\alpha,\beta}(\Gamma) \text{ if and only if } (x,y) \in \Omega_{\sigma\circ\alpha,\sigma\circ\beta}(\Gamma).$$

Then $\varphi(s(e)) = s(\psi(e))$, $\varphi(r(e)) = r(\psi(e))$ for all $e \in \Omega_{\alpha,\beta}(\Gamma)$ and $\lambda_x = \lambda_x^\sigma$ where λ^σ is the family of Radon measures for $Q_{\sigma\circ\alpha,\sigma\circ\beta}(\Gamma)$.

For the second isomorphism, use $\varphi = \sigma^{-1}$ and $\psi(x,y) = (\sigma^{-1}(x), \sigma^{-1}(y))$. Then $(x,y) \in r^{-1}(x)$ if and only if $\alpha(y) = \beta(x)$. Let $x_0 = \sigma^{-1}(x)$ and $y_0 = \sigma^{-1}(y)$, then $(x_0, y_0) \in r_\sigma^{-1}(\varphi(x))$ if and only if $(x,y) \in r^{-1}(x)$ where r_σ is the range map of $Q_{\alpha\circ\sigma,\beta\circ\sigma}(\Gamma)$. Furthermore, if $(x,y) \neq (x,y')$ in $\Omega_{\alpha,\beta}(\Gamma)$, then $y_0 = \sigma^{-1}(y) \neq \sigma^{-1}(y') = y_0'$ and so, $(x_0, y_0) \neq (x_0, y_0')$ in $\Omega_{\alpha\circ\sigma,\beta\circ\sigma}(\Gamma)$. Thus, we have shown

$$\lambda^\sigma_{\varphi(x)} \circ \psi = c\lambda_x \text{ for some } c > 0$$

where λ^σ is the family of Radon measure for $Q_{\alpha\circ\sigma,\beta\circ\sigma}(\Gamma)$ since both are left Haar measures on the same set.

Also, for $e = (x,y) \in \Omega_{\alpha,\beta}(\Gamma)$

$$\varphi(s(e)) = \sigma^{-1}(y) = s_\sigma(\psi(e)) \qquad \text{and} \qquad \varphi(r(e)) = \sigma^{-1}(x) = r_\sigma(\psi(e)).$$

□

Example 3.9. [24] Let α be an endomorphism of a unital C^*-algebra B. A *transfer operator* L for (B, α) is a positive linear map $L : B \to B$ such that $L(\alpha(a)b) = aL(b)$ for all $a, b \in B$. [24] calls the triple (B, α, L) an Exel system and create the Cuntz-Pimsner algebra out of the right Hilbert module M_L, the completion of B with respect to the left and right actions

$$a \cdot \xi \cdot b = a\xi\alpha(b)$$

and inner product

$$\langle \xi, \eta \rangle = L(\xi^*\eta).$$

When $B = C(\Gamma)$ for compact group Γ and $\alpha = \beta^\#$ where $\beta \in \operatorname{End}(\Gamma)$, we get

$$L(\xi)(x) = \lambda_x(\xi)$$

defines a transfer operator for $(C(\Gamma), \alpha)$ where λ is the family of Radon measures for $Q_{\beta,1}(\Gamma)$. First note that $\Omega_{\beta,1}(\Gamma) = \{(x,y) \mid x = \beta(y)\}$ and so $C(\Omega_{\beta,1}(\Gamma)) \cong C(\Gamma)$ where the left and right actions become

$$a \cdot \xi \cdot b = a\xi\beta^\#(b)$$

and inner product

$$\langle \xi, \eta \rangle(x) = \int_{s(r^{-1}(x))} \overline{\xi(y)}\eta(y)\, d\lambda_x(x,y) = L(\xi^*\eta)(x)$$

for $a, b \in C(\Gamma)$, $\xi, \eta \in M_L = \mathcal{E}_{\beta,1}(\Gamma)$ and $x \in \Gamma$. Now for $x \in \Gamma$, $a \in C(\Gamma)$ and $\xi \in M_L$,

$$\begin{aligned} L(\alpha(a)\xi)(x) &= \lambda_x(\alpha(a)\xi) \\ &= \int a(\beta(y))\xi(y)\, d\lambda_x(x,y) \\ &= \int a(x)\xi(y)\, d\lambda_x(x,y) \\ &= a(x)\lambda_x(\xi) \\ &= [aL(\xi)](x). \end{aligned}$$

This shows the C^*-algebras $\mathcal{O}_{\beta,1}(\Gamma)$ were also studied in [24]. Moreover, for the C^*-algebras $\mathcal{O}_{\alpha,\beta}(\Gamma)$, notice

$$\begin{aligned} \lambda_x(\alpha^{\#}(a)\cdot\xi) &= \int a(\alpha(s(e)))\xi(e)\, d\lambda_x(e) \\ &= \int a(\beta(r(e)))\xi(e)\, d\lambda_x(e) \\ &= \int a(\beta(x))\xi(e)\, d\lambda_x(e) \\ &= \beta^{\#}(a)(x)\lambda_x(\xi) \end{aligned}$$

and hence, these topological group relations extend the context in [24]

However, there are topological group relations that are not of this form; for instance, $\mathcal{O}_{1,0}(\mathbb{Z}_3) \ncong \mathcal{O}_{n,1}(\mathbb{Z}_3)$ for any $n = 0, 1, 2$ (see Example 3.12.)

Example 3.10. Let $\alpha \in \mathrm{End}\ (\Gamma)$ be surjective for some compact group Γ. Then

$$\Omega_{1,\alpha}(\Gamma) = \{(x, \alpha(x)) \,\big|\, x \in \Gamma\} \cong \Gamma.$$

Hence, the left and right actions become

$$(a \cdot \xi \cdot b)(x, \alpha(x)) = a(\alpha(x))\xi(x, \alpha(x))b(x)$$

and the inner product becomes

$$\langle \xi, \eta \rangle(x) = \overline{\xi(x, \alpha(x))}\eta(x, \alpha(x))$$

for $\xi, \eta \in C(\Omega_{1,\alpha}(\Gamma))$, $a, b \in C(\Gamma)$ and $x \in \Gamma$. We quickly note that, for ϕ the left action of $C(\Gamma)$, $\ker \phi = \{0\}$ since α is surjective and so $J_{\mathcal{E}_{1,\alpha}(\Gamma)} = J(\mathcal{E}_{1,\alpha}(\Gamma)) = \phi^{-1}(\mathcal{K}(C(\Omega_{1,\alpha}(\Gamma))))$. We also note, for any $a \in C(\Gamma)$, $\xi(x,y) = a(y)$ defines $\xi \in C(\Omega_{1,\alpha}(\Gamma))$ and so

$$\theta_{\xi,1}(\eta)(x, \alpha(x)) = \xi(x, \alpha(x))\langle 1, \eta \rangle(x) = a(\alpha(x))\eta(x, \alpha(x)) = a \cdot \eta(x, \alpha(x))$$

for all $(x, \alpha(x)) \in \Omega_{1,\alpha}(\Gamma)$ and $\eta \in C(\Omega_{1,\alpha}(\Gamma))$; that is,

$$J_{\mathcal{E}_{1,\alpha}(\Gamma)} = C(\Gamma).$$

Now we let $u \in C(\Gamma)$ be defined by $u(x, \alpha(x)) = 1$ for all $(x, \alpha(x)) \in \Omega_{1,\alpha}(\Gamma)$. Then note

$$[a \cdot u](x, \alpha(x)) = a(\alpha(x)) = [u \cdot \alpha^{\#}(a)](x, \alpha(x)),$$

$$\langle u, u\rangle = 1 \qquad \text{and} \qquad u\langle u, \xi\rangle = \xi$$

where $\alpha^{\#}(a)(x) = a(\alpha(x))$. Hence, by Theorem 2.18 with $S = S_u$,

$$\mathcal{O}_{1,\alpha}(\Gamma) \cong C^*(C(\Gamma), S \,|\, S^*S = SS^* = 1 \text{ and } \alpha^{\#}(a) = S^*aS \text{ for all } a \in C(\Gamma)).$$

That is, if $\alpha^{\#}$ is an automorphism, then

$$\mathcal{O}_{1,\alpha}(\Gamma) \cong C(\Gamma) \rtimes_{\alpha^{\#}} \mathbb{Z}$$

as (indirectly) noted by Pimsner [60] and in particular, if $\alpha \in \mathrm{Aut}(\Gamma)$ then

$$\mathcal{O}_{1,\alpha}(\Gamma) \cong C(\Gamma) \rtimes_{\alpha^{\#}} \mathbb{Z} \qquad \text{and} \qquad \mathcal{O}_{\alpha,1}(\Gamma) \cong \mathcal{O}_{1,\alpha^{-1}}(\Gamma) \cong C(\Gamma) \rtimes_{(\alpha^{-1})^{\#}} \mathbb{Z}.$$

3.2. Topological $\mathbb{Z}_p$−Quivers.

Example 3.11. Let Γ be the compact abelian group $\mathbb{Z}_p = \mathbb{Z}/p\mathbb{Z}$. Then $\mathcal{O}_{\alpha,\beta}(\Gamma)$ is, in general, a graph C^*-algebra, but this does not yield all such since $(1_\Gamma, 1_\Gamma) \in \Omega_{\alpha,\beta}(\Gamma)$ for all α and β. Recall $\mathrm{End}(\mathbb{Z}_p) \cong \mathbb{Z}_p$.



FIGURE 1. The $\mathbb{Z}_2$-quivers: From left to right: $Q_{0,0}(\mathbb{Z}_2)$, $Q_{0,1}(\mathbb{Z}_2)$, $Q_{1,0}(\mathbb{Z}_2)$, $Q_{1,1}(\mathbb{Z}_2)$



FIGURE 2. The $\mathbb{Z}_3$-quivers: From left to right and top to bottom: $Q_{0,0}(\mathbb{Z}_3)$, $Q_{0,1}(\mathbb{Z}_3) \cong Q_{0,2}(\mathbb{Z}_3)$, $Q_{1,0}(\mathbb{Z}_3) \cong Q_{2,0}(\mathbb{Z}_3)$, $Q_{1,1}(\mathbb{Z}_3) \cong Q_{2,2}(\mathbb{Z}_3)$, $Q_{1,2}(\mathbb{Z}_3) \cong Q_{2,1}(\mathbb{Z}_3)$

Example 3.12. It can be shown that the graph C^*-algebra of a graph with a single loop on one vertex is $\mathcal{O}_1 \cong C(\mathbb{T})$. Hence, $\mathcal{O}_{1,1}(\mathbb{Z}_2) \cong C(\mathbb{T}) \oplus C(\mathbb{T})$, and $\mathcal{O}_{1,1}(\mathbb{Z}_3)) \cong C(\mathbb{T})^{\oplus 3}$. Furthermore, it is shown in [29] that $\mathcal{O}_{0,1}(\mathbb{Z}_3) \cong C(S^2_{q,1})$ for any $q \in \mathbb{R}\backslash[0,1]$ (see [29] for further details). Finally, the author shows $\mathcal{O}_{1,0}(\mathbb{Z}_2) \cong \mathcal{T}$ and $\mathcal{O}_{1,2}(\mathbb{Z}_3) = C(\mathbb{T}) \oplus M_2(C(\mathbb{T}))$ in [51].

Just a few interesting facts concerning these types of topological group quivers follow:

Proposition 3.13. Let $n \in \mathbb{Z}_p$ be non-zero. Then

$$Q_{n,n}(\mathbb{Z}_p) \cong \sqcup_{j=1}^{p/\gcd(n,p)} Q_{0,0}(\mathbb{Z}_{\gcd(n,p)}).$$

Moreover,

$$\mathcal{O}_{n,n}(\mathbb{Z}_p) \cong \oplus_{j=1}^{p/\gcd(n,p)} \mathcal{O}_{0,0}(\mathbb{Z}_{\gcd(n,p)}).$$

Proof. Let $z = e^{2\pi i/p}$ and let $n_0 = \gcd(n,p)$. Denote $Z_k = \{e^{2(k+mp/n_0)\pi i/p} \,\big|\, m = 0, ..., n_0 - 1\}$ for $k = 0, ..., p/n_0 - 1$. Then note for any $x, y \in Z_k$ where $x = e^{2(k+m_1 p/n_0)\pi i/p}$ and $y = e^{2(k+m_2 p/n_0)\pi i/p}$,

$$(xy^{-1})^{n_0} = e^{2(m_1-m_2)\pi i} = 1;$$

that is, $x^{n_0} = y^{n_0}$. Furthermore, if $x = e^{2(k_1+m_1p/n_0)\pi i/p} \in Z_{k_1}$ and $y = e^{2(k_2+m_2p/n_0)\pi i/p} \in Z_{k_2}$ for $k_1 \neq k_2$ then

$$(xy^{-1})^{n_0} = e^{2(k_1-k_2)n_0\pi i/p + 2(m_1-m_2)\pi i} = e^{2(k_1-k_2)n_0\pi i/p}$$

but $|k_1 - k_2| < p/n_0$ ensuring that $(xy^{-1})^{n_0} \neq 1$ (or better written, $x^{n_0} \neq y^{n_0}$.) The rest is now a trivial application of Theorem 2.22.

□

It can be shown (see [43], [23], [1]) that the graph with m-points connected in a single length m loop admits the universal C^*-algebra $M_m(C(\mathbb{T}))$. Let $\mathbb{Z}_p^*$ denote the multiplicative group $\{n \in \mathbb{Z}_p \,\big|\, \gcd(n,p) = 1\}$.

Proposition 3.14. Let $n, p \in \mathbb{N}$ such that $\gcd(n,p) = 1$. Then:

1. $Q_{n,1}(\mathbb{Z}_p)$ consists of disjoint loops of various lengths less than or equal to $o_p(n)$ (the order of n in $\mathbb{Z}_p^*$).
2. $Q_{1,n}(\mathbb{Z}_p) \cong Q_{n,1}(\mathbb{Z}_p)$.
3. The number of base points of loops of length $k \leq o_p(n)$ in $Q_{n,1}(\mathbb{Z}_p)$ is
$$\gcd(n^k - 1, p).$$

Proof. (1) Given $z \in \mathbb{Z}_p$, there exists a unique edge with range z, likewise source z, since $\gcd(n,p) = \gcd(1,p) = 1$. Furthermore, there exists a number $k \leq o_p(n)$ such that $z^{n^k} = z$.

(2) It is clear from (1) that reversing the arrows will not change the graph ($Q_{1,n}(\mathbb{Z}_p)$ is $Q_{n,1}(\mathbb{Z}_p)$ with reversed arrows).

(3) We need only find all solutions, y, to $n^k y = y \mod p$. This equation has $\gcd(n^k - 1, p)$ distinct solutions.

□

Example 3.15. With this proposition, it is easy to determine the C^*-algebra $\mathcal{O}_{n,1}(\mathbb{Z}_p)$ where $\gcd(n,p) = 1$.

(1) $\mathcal{O}_{5,1}(\mathbb{Z}_{72})$: Note $o_{72}(5) = 6$ and

$$\begin{aligned}
\gcd(5-1, 72) &= \gcd(4, 72) = 4\\
\gcd(5^2-1, 72) &= \gcd(24, 72) = 24\\
\gcd(5^3-1, 72) &= \gcd(124, 72) = 4\\
\gcd(5^4-1, 72) &= \gcd(624, 72) = 24\\
\gcd(5^5-1, 72) &= \gcd(3124, 72) = 4\\
\gcd(5^6-1, 72) &= \gcd(15624, 72) = 72.
\end{aligned}$$

Hence, there are 4 base points of 1-loops and 24 base points of 2-loops. Note: each 1-loop counts here, and so there are 20 base points of 2-loops that aren't base points of 1-loops; that is, 10 loops of length 2. Furthermore, the 4 base points of 3-loops are already counted by the 1-loops, and so on... Finally, there are 72 base points of 6-loops. Subtracting all possible base points of 2-loops, we obtain 48 base points of 6-loops and hence, 8 6-loops. Therefore,

$$\mathcal{O}_{5,1}(\mathbb{Z}_{72}) = C(\mathbb{T})^{\oplus 4} \oplus M_2(C(\mathbb{T}))^{\oplus 10} \oplus M_6(C(\mathbb{T}))^{\oplus 8}.$$

Remark: $4 + 2(10) + 6(8) = 72$.

(2) $\mathcal{O}_{6,1}(\mathbb{Z}_{77})$: Note $o_{77}(6) = 10$ and

$$\gcd(6-1, 77) = 1 \qquad \gcd(6^2-1, 77) = 7 \qquad \gcd(6^{10}-1, 77) = 77.$$

That is, there is one copy of $C(\mathbb{T})$, $\frac{7-1}{2} = 3$ copies of $M_2(C(\mathbb{T}))$, and $\frac{77-7}{10} = 7$ copies of $M_{10}(C(\mathbb{T}))$; i.e.,

$$\mathcal{O}_{6,1}(\mathbb{Z}_{77}) = C(\mathbb{T}) \oplus M_2(C(\mathbb{T}))^{\oplus 3} \oplus M_{10}(C(\mathbb{T}))^{\oplus 7}.$$

Remark: $1 + 2(3) + 10(7) = 77$.

Proposition 3.16. Let p be prime and let $\varphi(p-1)$ denote the number of divisors of $p-1$. Then the number of distinct (non-isomorphic) topological $\mathbb{Z}_p$-quivers is $\varphi(p-1)+3$.

Proof. Begin by noting that $\mathrm{Aut}(\mathbb{Z}_p) = \mathbb{Z}_p^* = \{1, ..., p-1\}$, the numbers coprime to the prime number p. Hence, by Proposition 3.8,

$$Q_{n,m}(\mathbb{Z}_p) \cong Q_{kn,km}(\mathbb{Z}_p)$$

for each $k = 1, ..., p-1$. This gives us only the following quivers to check

$$Q_{0,0}(\mathbb{Z}_p), Q_{0,1}(\mathbb{Z}_p), Q_{1,0}(\mathbb{Z}_p) \text{ [the 3 in } \varphi(p-1)+3\text{], and } Q_{1,n}(\mathbb{Z}_p)$$

where $n = 1, ..., p-1$. We see immediately that $Q_{0,0}(\mathbb{Z}_p), Q_{0,1}(\mathbb{Z}_p)$ and $Q_{1,0}(\mathbb{Z}_p)$ are all distinct since $\Omega_{0,0}(\mathbb{Z}_p) = \mathbb{Z}_p \times \mathbb{Z}_p$, $\Omega_{0,1}(\mathbb{Z}_p) = \{1\} \times \mathbb{Z}_p$ (i.e., all arrows have range equal 1) and $\Omega_{1,0}(\mathbb{Z}_p) = \mathbb{Z}_p \times \{1\}$ (i.e., all arrows have source equal 1.) Furthermore, $Q_{1,n}(\mathbb{Z}_p)$ has one and only one arrow with range k (likewise, with source k) so $Q_{1,n}(\mathbb{Z}_p)$ is not isomorphic to any of $Q_{0,0}(\mathbb{Z}_p), Q_{0,1}(\mathbb{Z}_p)$, or $Q_{1,0}(\mathbb{Z}_p)$. In fact, $Q_{1,n}(\mathbb{Z}_p)$ is solely defined based on the length of its loops; that is, the order of n in $\mathbb{Z}_p^*$ as the previous proposition proves. Since all orders divide $p-1$ and all orders are achieved, the resulting count of distinct topological $\mathbb{Z}_p$-quivers is $\varphi(p-1)+3$. □

3.3. Topological $\mathbb{T}^d$-Quivers.

Example 3.17. For the compact abelian group $\mathbb{T}^d$, note $\mathrm{End}(\mathbb{T}^d) \cong M_d(\mathbb{Z})$ ([67]); that is, an element $\sigma \in \mathrm{End}(\mathbb{T}^d)$ is of the form σ_F for some $F \in M_d(\mathbb{Z})$ where

$$\sigma_F(e^{2\pi i t}) = e^{2\pi i F t}$$

for each $t \in \mathbb{Z}^d$. To simplify notation, use F and G in place of σ_F and σ_G whenever convenient. For instance,

$$Q_{F,G}(\mathbb{T}^d) := Q_{\sigma_F, \sigma_G}(\mathbb{T}^d)$$

and the C^*-correspondence

$$\mathcal{E}_{F,G}(\mathbb{T}^d) := \mathcal{E}_{\sigma_F,\sigma_G}(\mathbb{T}^d)$$

where $F, G \in M_d(\mathbb{Z})$. We will consider the cases when these maps are surjective; that is, $\det F$ and $\det G$ are non-zero.

Let $F, G \in M_d(\mathbb{Z})$ where $\det F, \det G \neq 0$. Then $|\ker \sigma_F| = |\det F|$ and so, the $C(\mathbb{T}^d)$-valued inner product becomes

$$\langle \xi, \eta \rangle(x) = \frac{1}{|\det F|} \sum_{\sigma_F(y)=\sigma_G(x)} \overline{\xi(x,y)}\eta(x,y)$$

for $\xi, \eta \in \mathcal{E}_{F,G}(\mathbb{T}^d) = C(\Omega_{F,G}(\mathbb{T}^d))$ [by Lemma 3.6] and $x \in \mathbb{T}^d$. This is a finite sum since the number of solutions, y, to $\sigma_F(y) = \sigma_G(x)$ given any $x \in \mathbb{T}^d$ is $|\det F| < \infty$.

Remark 3.18. The left action, ϕ, is defined by

$$\phi(a)\xi(x,y) = a(y)\xi(x,y)$$

for $a \in C(\mathbb{T}^d)$, $\xi \in C(\Omega_{F,G}(\mathbb{T}^d))$ and $(x,y) \in \Omega_{F,G}(\mathbb{T}^d)$. Note: ϕ is injective. To see this claim, let $a \in C(\mathbb{T}^d)$ and assume $\phi(a)\xi = 0$ for each $\xi \in C(\Omega_{F,G}(\mathbb{T}^d))$. Then $a(y)\xi(x,y) = 0$ for each $(x,y) \in \Omega_{F,G}(\mathbb{T}^d)$ and $\xi \in C(\Omega_{F,G}(\mathbb{T}^d))$. Since $s(\Omega_{F,G}(\mathbb{T}^d)) = \{y \in \mathbb{T}^d \,|\, (x,y) \in \Omega_{F,G}(\mathbb{T}^d)\} = \mathbb{T}^d$ by the surjectivity of σ_F, $a = 0$.

Examples of the form $G = 1_d$ are studied in [24] and examples when $d = 1$ can be found in [68] where $\gcd(n,m) = 1$. Brenken [7] examines cases when $F = a1_d$ and G is arbitary, and examines the associated C^*-algebra when $d = 1$, $F = a \in \mathbb{N}$ and $G = 1$.

In fact, a basis for $\mathcal{E}_{F,G}(\mathbb{T}^d)$ can be provided using the following lemma and corollaries to Proposition 3.8.

Proposition 3.19. If $U, F, G \in M_d(\mathbb{Z})$ such that $\det U = 1$ and $\det F, \det G \neq 0$, then $\mathcal{O}_{F,G}(\mathbb{T}^d) \cong \mathcal{O}_{UF,UG}(\mathbb{T}^d) \cong \mathcal{O}_{FU,GU}(\mathbb{T}^d)$.

Proof. This is evident from Proposition 3.8. □

Corollary 3.20. Given $F, G \in M_d(\mathbb{Z})$ such that $\det F, \det G \neq 0$, there exist $F', G' \in M_d(\mathbb{Z})$ with F' a positive diagonal matrix and

$$\mathcal{O}_{F,G}(\mathbb{T}^d) \cong \mathcal{O}_{F',G'}(\mathbb{T}^d).$$

Proof. Using the Smith normal form (see [22]), there exist unimodular matrices $U, V \in M_d(\mathbb{Z})$ and a positive diagonal matrix $D \in M_d(\mathbb{Z})$ such that

$$F = UDV.$$

Hence $U^{-1}, V^{-1} \in M_d(\mathbb{Z})$ are unimodular matrices and, by the previous proposition,

$$\mathcal{O}_{F,G}(\mathbb{T}^d) \cong \mathcal{O}_{D,U^{-1}GV^{-1}}(\mathbb{T}^d).$$

□

The following lemma is well known:

Lemma 3.21. Given Γ_0 a finite subgroup of $\mathbb{T}^d$, $\sum_{z \in \Gamma_0} z = 0$ (in $\mathbb{C}^d$.)

Let $F = \operatorname{Diag}(a_1, ..., a_d) \in M_d(\mathbb{Z}), G = (b_{jk})_{j,k=1}^d \in M_d(\mathbb{Z})$ where $a_j > 0$ for each $j = 1, ..., d$, $\det G \neq 0$ and let G_j denote the j-th row of G, $(b_{jk})_{k=1}^d$. Further, let $N = \det F = \prod_{j=1}^d a_j > 0$ and let

$$\mathfrak{I}(F) = \{\nu = (\nu_j)_{j=1}^d \in \mathbb{Z}^d \,\big|\, 0 \leq \nu_j \leq a_j - 1\}.$$

The $C(\mathbb{T}^d)$-valued inner product becomes

$$\langle \xi, \eta \rangle(x) = \frac{1}{N} \sum_{\sigma_F(y)=\sigma_G(x)} \overline{\xi(x,y)}\eta(x,y)$$

for all $\xi, \eta \in C(\Omega_{F,G}(\mathbb{T}^d))$ and $x \in \mathbb{T}^d$.

Given $\nu \in \mathfrak{I}(F)$, define $u_\nu \in C(\Omega_{F,G}(\mathbb{T}^d))$ by

$$u_\nu(x,y) = y^\nu = \prod_{j=1}^d y^{\nu_j}$$

for $(x,y) \in \Omega_{F,G}(\mathbb{T}^d)$. So $u_{(0,0,...,0)} = 1 \in C(\Omega_{F,G}(\mathbb{T}^d))$. Then

$$\begin{aligned}
\langle u_\nu, u_{\nu'} \rangle(x) &= \frac{1}{N} \sum_{\sigma_F(y)=\sigma_G(x)} \overline{y}^\nu y^{\nu'} \\
&= \frac{1}{N} \sum_{z \in \ker \sigma_F} z_0^{\nu-\nu'} z^{\nu-\nu'} \qquad \text{for any } z_0 \in \sigma_F^{-1}(\sigma_G(x)) \\
&= \begin{cases} 0 & \text{if } \nu \neq \nu' \\ 1 & \text{if } \nu = \nu' \end{cases} = \delta_\nu^{\nu'}.
\end{aligned}$$

Furthermore, since

$$u_\nu \langle u_\nu, \xi \rangle(x,y) = u_\nu(x,y)\langle u_\nu, \xi \rangle(x) = y^\nu \langle u_\nu, \xi \rangle(x),$$

we have

$$\begin{aligned}
[\sum_{\nu \in \mathfrak{I}(F)} u_\nu \langle u_\nu, \xi \rangle](x,y) &= \frac{1}{N} \sum_{\nu \in \mathfrak{I}(F)} \sum_{\sigma_F(w)=\sigma_G(x)} y^\nu \overline{w}^\nu \xi(x,w) \\
&= \frac{1}{N} \sum_{\nu} \sum_{\sigma_F(w)=\sigma_G(x)} [\prod_{j=1}^d y_j^{\nu_j} \overline{w}_j^{\nu_j}]\xi(x,w).
\end{aligned}$$

Note: if $x = e^{2\pi i t}$ and $\sigma_F(w) = \sigma_G(x)$, then $w_j^{a_j} = e^{2\pi i (Gt)_j}$ where $(Gt)_j$ denotes the j-th coordinate of the vector Gt. Now note

$$\sum_{k=0}^{a_j-1} y_j^k \overline{w}_j^k = \begin{cases} \frac{(y_j\overline{w}_j)^{a_j}-1}{y_j\overline{w}_j-1} & \text{if } y_j \neq w_j \\ a_j & \text{if } y_j = w_j \end{cases} = \begin{cases} 0 & \text{if } x_j \neq w_j \\ a_j & \text{if } x_j = w_j \end{cases}$$

since $y_j^{a_j}\overline{w}_j^{a_j} = e^{2\pi i(Gt)_j}e^{-2\pi i(Gt)_j} = 1$ where $x = e^{2\pi it}$ $(t \in \mathbb{Z}^d)$. Hence,

$$[\sum_{\nu\in\mathfrak{I}(F)} u_\nu\langle u_\nu, \xi\rangle](x,y) = \frac{1}{N}\sum_{\nu}\sum_{\sigma_F(w)=\sigma_G(x)}[\prod_{j=1}^{d} y_j^{\nu_j}\overline{w}_j^{\nu_j}]\xi(x,w)$$

$$= \frac{1}{N}[\prod_{j=1}^{d} a_j]\xi(x,y) = \xi(x,y).$$

Thus, $\{u_\nu\}_{\nu\in\mathfrak{I}(F)}$ is a basis for $\mathcal{E}_{F,G}(\mathbb{T}^d)$.

For $\nu \in \mathfrak{I}(F)$ and $j \in \{1,...,d\}$, using notation from Theorem 2.18, set $S_\nu = S_{u_\nu}$, $S = S_{(0,0,...,0)}$ and let $\{U_j\}_{j=1}^d$ be the full spectrum unitaries in $C(\mathbb{T}^d)$ defined by $U_j(y) = y_j$ for $y \in \mathbb{T}^d$. Then, using the notation

$$U^\nu = U_1^{\nu_1}U_2^{\nu_2}\cdots U_d^{\nu_d},$$

we have

$$U^\nu S = S_{U^\nu\cdot u_0} = S_\nu$$

since $[U^\nu \cdot u_0](x,y) = y^\nu = u_\nu(x,y)$, and

$$U_j^{a_j}S = S_{U_j^{a_j}\cdot u_0} = S\prod_{k=1}^{d} U_k^{b_{jk}} = SU^{G_j}$$

since $[U_j^{a_j} \cdot u_0](x,y) = y_j^{a_j} = u_0 \cdot U_j \circ \sigma_G(x) = u_0 \cdot \prod_{k=1}^d U_k^{b_{jk}}(x,y) = u_0 \cdot U^{G_j}(x,y)$.

Remark 3.22. [53, Proposition 2.21] states that if the left action ϕ is injective then the natural inclusion $C(\mathbb{T}^d) \hookrightarrow \mathcal{O}_{F,G}(\mathbb{T}^d)$ is injective. Hence, the image of U_j in $\mathcal{O}_{F,G}(\mathbb{T}^d)$ (still denoted U_j) is a full spectrum unitary since U_j is a full spectrum unitary in $C(\mathbb{T}^d)$.

Thus, by Theorem 2.18:

Theorem 3.23. *Let $F = \mathrm{Diag}(a_1,...,a_d), G \in M_d(\mathbb{Z})$ where $\det F, \det G \neq 0$ and let G_j be the j-th row vector of G. Further, let $\mathfrak{I}(F)$ denote the set $\{\nu = (\nu_j)_{j=1}^d \in \mathbb{Z}^d \mid 0 \leq \nu_j \leq a_j - 1\}$. Then $\mathcal{O}_{F,G}(\mathbb{T}^d)$ is the universal C^*-algebra generated by isometries $\{S_\nu\}_{\nu\in\mathfrak{I}(F)}$ and (full spectrum) commuting unitaries $\{U_j\}_{j=1}^d$ that satisfy the relations*

1. $S_\nu^*S_{\nu'} = \langle u_\nu, u_{\nu'}\rangle = \delta_\nu^{\nu'}$,
2. $U^\nu S = S_\nu$ *for all* $\nu \in \mathfrak{I}(F)$,
3. $U_j^{a_j}S = SU^{G_j}$, *for all* $j = 1,...,d$ *and*
4. $1 = \sum_{\nu\in\mathfrak{I}(F)} S_\nu S_\nu^* = \sum_{\nu\in\mathfrak{I}(F)} U^\nu SS^*U^{-\nu}$

where U^ν denotes $\prod_{j=1}^d U_j^{\nu_j}$.

Corollary 3.24. *Let $n \in \mathbb{N}$ and $G \in M_d(\mathbb{Z})$ where $\det G \neq 0$, then $\mathcal{O}_{n,G}(\mathbb{T}^d) := \mathcal{O}_{n1_d,G}(\mathbb{T}^d)$ is the universal C^*-algebra*

$$C^*(\{U_j\}_{j=1}^d, S \mid U_j^*U_j = U_jU_j^* = \sum_{\nu\in\mathfrak{n}^d} U^\nu SS^*U^{-\nu} = 1, S^*U^\mu S = \delta_0^\mu\ (\mu \in \mathfrak{n}^d), U_j^nS = SU^{G_j})$$

where $\mathfrak{n}^d = \{0,...,n-1\}^d$.

Corollary 3.25. Let $n \in \mathbb{N}$ and $m \in \mathbb{Z} \setminus \{0\}$, then $\mathcal{O}_{n,m}(\mathbb{T})$ is the universal C^*-algebra

$$C^*(U, S \,\big|\, U^*U = UU^* = \sum_{k=0}^{n-1} U^k SS^* U^{-k} = 1, S^*U^kS = \delta_0^k \ (0 \leq k \leq n-1), U^nS = SU^m).$$

3.4. **A Topological $\mathbb{T}^\omega$-Quiver and Some Generalizations.** We describe an example with a perfect compact $\ker \alpha$ which is $\mathbb{T}$. Let $\Gamma = \mathbb{T}^\omega := \prod_{n\in\mathbb{N}} \mathbb{T}$ be the compact abelian topological group with the product topology. Tychonoff's Theorem guarantees that Γ is a compact group. Let $\sigma \in \operatorname{End}(\mathbb{T}^\omega)$ be the (surjective) group endomorphism

$$\sigma(t_1, t_2, t_3, ...) = (t_2, t_3, ...)$$

for each $(t_1, t_2, t_3, ...) \in \mathbb{T}^\omega$. Then

$$\Omega_{\sigma,1}(\mathbb{T}^\omega) = \{(x, y) \in \mathbb{T}^\omega \times \mathbb{T}^\omega \,\big|\, \sigma(y) = x\},$$

$$r^{-1}(x) = \{x\} \times \{(z, x) \,\big|\, z \in \mathbb{T}\},$$

where $(z, x) = (z, x_1, x_2, ...) \in \mathbb{T}^\omega$, and finally, λ_x is the Haar measure on $\mathbb{T}$. The $C(\mathbb{T}^\omega)$-valued inner product becomes

$$\langle \xi, \eta \rangle(x) = \int_{\mathbb{T}} \overline{\xi(x, (z, x))} \eta(x, (z, x))\, dz$$

for each $x \in \mathbb{T}^\omega$.

Also note, given $a \in C(\mathbb{T}^\omega)$ and $\xi \in C(\Omega_{\sigma,1}(\mathbb{T}^\omega))$,

$$\sigma^\#(a) \cdot \xi(x, y) = a(\sigma(y))\xi(x, y) = a(x)\xi(x, y) = (\xi \cdot a)(x, y)$$

for each $(x, y) \in \Omega_{\sigma,1}(\mathbb{T}^\omega)$. Hence,

$$\sigma^\#(a)S_\xi = S_{\sigma^\#(a)\cdot\xi} = S_{\xi\cdot a} = S_\xi a$$

for each $a \in C(\mathbb{T}^\omega)$ and $\xi \in C(\Omega_{\sigma,1}(\mathbb{T}^\omega))$ where S_ξ is the image of ξ into the universal C^*-algebra $\mathcal{O}_{\sigma,1}(\mathbb{T}^\omega)$.

This result is, in fact, a special case of the following:

Proposition 3.26. Let Γ be a locally compact group and $\alpha, \beta \in \operatorname{End}(\Gamma)$. For any $\xi \in C(\Omega_{\alpha,\beta}(\Gamma))$ and $a \in C_0(\Gamma)$, we get

$$\pi(\alpha^\#(a))T(\xi) = T(\xi)\pi(\beta^\#(a))$$

where (T, π) is the universal representation coisometric on $J_{\mathcal{E}_{Q_{\alpha,\beta}(\Gamma)}}$ onto $\mathcal{O}_{\alpha,\beta}(\Gamma)$.

Proof. Let $a \in C_0(\Gamma)$ and $\xi \in C(\Omega_{\alpha,\beta}(\Gamma))$ then

$$\begin{aligned}(\alpha^\#(a) \cdot \xi)(x, y) &= \alpha^\#(a)(y)\xi(x, y)\\ &= a(\alpha(y))\xi(x, y)\\ &= \xi(x, y)a(\beta(x))\\ &= (\xi \cdot \beta^\#(a))(x, y)\end{aligned}$$

for each $(x, y) \in \Omega_{\alpha,\beta}(\Gamma)$ and hence,

$$\pi(\alpha^{\#}(a))T(\xi) = T(\alpha^{\#}(a) \cdot \xi) = T(\xi \cdot \beta^{\#}(a)) = T(\xi)\pi(\beta^{\#}(b)).$$

□

Theorem 3.27. Suppose Γ is a compact group with $\alpha, \beta \in \mathrm{End}(\Gamma)$ and the left action ϕ is injective (i.e., $Q_{\alpha,\beta}(\Gamma)$ is sinkless). If there exists an orthonormal basis $\{u_i\}_{i=1}^n$ then $\mathcal{O}_{\alpha,\beta}(\Gamma)$ is the universal C^*-algebra generated by $A = C(\Gamma)$ and isometries $\{S_i\}_{i=1}^n$ subject to

1. $\alpha^{\#}(a)S_i = S_i\beta^{\#}(a)$ for each $i = 1, ..., n$ and $a \in C(\Gamma)$
2. $S_i^* S_j = \delta_i^j$
3. $\sum_{i=1}^n S_i S_i^* = 1$.

If, in addition, there exist $\{a_i\}_{i=1}^n \subseteq C(\Gamma)$ and $u \in \{u_i\}_{i=1}^n$ such that $a_i \cdot u = u_i$ for each $i = 1, ..., n$ then $\mathcal{O}_{\alpha,\beta}(\Gamma)$ is the universal C^*-algebra generated by $A = C(\Gamma)$ and an isometry S subject to

1. $\alpha^{\#}(a)S = S\beta^{\#}(a)$ for each $i = 1, .., n$ and $a \in C(\Gamma)$
2. $S^* a_i S = \delta_0^i$
3. $\sum_{i=1}^n a_i S S^* a_i^* = 1$.

Proof. This is none other than a special case of Theorem 2.18.

□

Corollary 3.28. Let Γ be a compact group with $\alpha, \beta \in \mathrm{End}(\Gamma)$ and ϕ injective. If there exists $u \in \mathcal{E}_{\alpha,\beta}(\Gamma)$ where $u\langle u, \xi\rangle = \xi$ for each $\xi \in \mathcal{E}_{\alpha,\beta}(\Gamma)$ (i.e., $\theta_{u,u} = 1$) then $\mathcal{O}_{\alpha,\beta}(\Gamma)$ is the universal C^*-algebra generated by $A = C(\Gamma)$ and unitary S such that

$$\alpha^{\#}(a)S = S\beta^{\#}(a)$$

for all $a \in C(\Gamma)$.

Remark 3.29. This corollary suggests a similarity to crossed products by $\mathbb{Z}$. In some sense, what we have here is a "crossed product" by two endomorphisms. In fact, none of these last results are exclusive to topological group relations. Given a second countable locally compact Hausdorff space X, define a topological quiver $Q = (X, \Omega_{\alpha,\beta}(X), r, s, \lambda)$ with

$$\Omega_{\alpha,\beta}(X) = \{(x, y) \,|\, \alpha(y) = \beta(x)\}$$

for continuous maps α, β on X and appropriate measures λ. Then similar conclusions remain true.

4. Some C^*-subalgebras of $\mathcal{O}_{F,G}(\mathbb{T}^d)$

By Proposition 3.8, we may assume (for the remainder of this paper) that $F = \mathrm{Diag}(a_1, ..., a_d)$ with $a_j > 0$. Let $G \in M_d(\mathbb{Z})$ with j-th row denoted G_j and $\det G \neq 0$, then recall the definitions of $\{U_j\}_{j=1}^d$ and S from Section 3.3. $U_j \in C(\mathbb{T}^d)$ is the unitary defined by $U_j(x) = x_j$ for each $x = (x_k)_{k=1}^d \in \mathbb{T}^d$, $j \in \{1, ..., d\}$ and S is the isometry defined as the image of $u_0 = 1 \in C(\Omega_{F,G}(\mathbb{T}^d))$ into $\mathcal{O}_{F,G}(\mathbb{T}^d)$. By Theorem 3.23, the C^*-algebra $\mathcal{O}_{F,G}(\mathbb{T}^d)$ is the universal C^*-algebra generated by the commuting unitaries $\{U_j\}_{j=1}^d$ and isometry S satisfying:

(1) $S^*U^\nu S = \delta_0^\nu$, for all $\nu \in \mathfrak{I}(F)$,
(2) $U_j^{a_j} S = SU^{G_j}$, for all $j = 1, ..., d$ and
(3) $\sum_{\nu \in \mathfrak{I}(F)} U^\nu SS^* U^{-\nu} = 1$

where $\mathfrak{I}(F) = \{\nu = (\nu_j)_{j=1}^d \in \mathbb{Z}^d \,\big|\, 0 \le \nu_j \le a_j - 1\}$.

Lemma 4.1. Given $k \in \mathbb{N}$ and $|\det G| = 1$, the C^*-algebra $C^*(\{U_j\}_{j=1}^d, S^k)$ viewed as a C^*-subalgebra of $\mathcal{O}_{F,G}(\mathbb{T}^d)$ is a quotient of $\mathcal{O}_{F^k,G^k}(\mathbb{T}^d)$.

Proof. Let $\tilde{S}_\nu = U^\nu S^k$ for $\nu \in \mathfrak{I}(F^k) = \{(\nu_j)_{j=1}^d \,\big|\, 0 \le \nu_j \le a_j^k - 1\}$ ($\tilde{S} = S^k$) and so

(1) $\tilde{S}^* U^{\nu'} \tilde{S} = (S^k)^* U^{\nu'} S^k = \delta_0^{\nu'}$, for each $\nu' \in \mathfrak{I}(F^k)$.

Also, by (3) of Theorem 3.23,

$$U^{a_j G_j} S = [\prod_{m=1}^d U_m^{a_j b_{jm}}] S = S \prod_{m,l=1}^d U_l^{b_{jm} b_{ml}} = SU^{(G^2)_j}.$$

Thus,

2. $U_j^{a_j^k} \tilde{S} = SU^{a_j^{k-1} G_j} S^{k-1} = ... = S^k U^{(G^k)_j} = \tilde{S} U^{(G^k)_j}$

For $k = 2$, we may write $\nu = \nu_0 + \mu \cdot \nu_1$ [point-wise product and sum] where $\mu = (a_1, ..., a_d)$, $\nu_i \in \mathfrak{I}(F)$. Next, note for any $\omega \in \mathfrak{I}(F)$, there exist unique $\nu_1 \in \mathfrak{I}(F)$ and unique $k_{\nu_1} \in \mathbb{Z}^d$ such that $(\nu_1 + k_{\nu_1} \cdot \mu)G = \omega$ (matrix multiplication on the right) by the bijectivity of G. Hence,

$$\begin{aligned}\sum_{\nu_1 \in \mathfrak{I}(F)} U^{\nu_1 G} SS^* U^{-\nu_1 G} &= \sum_{\nu_1} U^{(\nu_1 + k_{\nu_1} \cdot \mu)G} U^{(k_{\nu_1} \cdot \mu)G} SS^* U^{-(k_{\nu_1} \cdot \mu)G} U^{-(\nu_1 + k_{\nu_1} \cdot \mu)G} \\ &= \sum_{\nu_1} U^{(\nu_1 + k_{\nu_1} \cdot \mu)G} SU^{k_{\nu_1} G^2} U^{-k_{\nu_1} G^2} S^* U^{-(\nu_1 + k_{\nu_1} \cdot \mu)G} \\ &= \sum_{\nu_1} U^{(\nu_1 + k_{\nu_1} \cdot \mu)G} SS^* U^{-(\nu_1 + k_{\nu_1} \cdot \mu)G} \\ &= \sum_{\nu_1' \in \mathfrak{I}(F)} U^{\nu_1' G} SS^* U^{-\nu_1' G} = 1.\end{aligned}$$

This gives us

$$\begin{aligned}\sum_{\nu \in \mathfrak{I}(F^k)} U^\nu \tilde{S} \tilde{S}^* U^{-\nu} &= \sum_{\nu_0, \nu_1 \in \mathfrak{I}(F)} U^{\nu_0} U^{\mu\nu_1} S^2 (S^2)^* U^{-\mu\nu_1} U^{-\nu_0} \\ &= \sum_{\nu_0, \nu_1} U^{\nu_0} S [U^{\sum_{j=1}^d (\nu_1)_j G_j} SS^* U^{-\sum_{j=1}^d (\nu_1)_j G_j}] S^* U^{-\nu_0} \\ &= \sum_{\nu_0} U^{\nu_0} SS^* U^{-\nu_0} = 1.\end{aligned}$$

A similar argument can be used to show $\sum_{\nu \in \mathfrak{I}(F^k)} \tilde{S}_\nu \tilde{S}_\nu^* = 1$ where $k \in \mathbb{N}$. Hence, by Theorem 3.23, universality implies that there exists a surjective homomorphism of $\mathcal{O}_{F^k,G^k}(\mathbb{T}^d)$ onto $C^*(\{U_j\}_{j=1}^d, S^k)$.

□

Lemma 4.2. If $k_j|a_j$ for each $j = 1, ..., d$ and $\gcd(\det F, \det G) = 1$ then

$$C^*(\{U_j^{k_j}\}_{j=1}^d, S) = \mathcal{O}_{F,G}(\mathbb{T}^d).$$

Next assume $k_j|a_j$ for all $j = 1, ..., d$. Then there are $p_j \in \mathbb{N}$ such that $a_j = k_j p_j$, hence

$$U_j^{a_j} = (U_j^{k_j})^{p_j} \in C^*(\{U_j^{k_j}\}_{j=1}^d, S)$$

for all $j = 1, ..., d$. Further note $U^{G_j} = S^* U_j^{a_j} S \in C^*(\{U_j^{k_j}\}_{j=1}^d, S)$.

Let $Q = (q_{ij})_{ij} = [Q_1 ... Q_d]^T = \text{adj } G \in M_d(\mathbb{Z})$, the adjugate of G and $G = [G'_1 ... G'_d]$ so that $QG = GQ = (\det G)1_d$ and hence, $Q_j G'_m = (\det G)\delta_j^m$. Then given $t \in \mathbb{T}^d$ and $j \in \{1, ..., d\}$, we have

$$\sum_{l=1}^d q_{jl} G_l t = \sum_{m=1}^d \sum_{l=1}^d (q_{jl} b_{lm}) t_m = \sum_m Q_j G'_m t_m = (\det G) t_j$$

showing that

$$U_j^{\det G} = \prod_l (U^{G_l})^{q_{jl}} \in C^*(\{U_j^{k_j}\}_{j=1}^d, S).$$

Finally, since $\gcd(\det F, \det G) = 1$, we have $\gcd(a_j, \det G) = 1$ for each j. Hence, there exists $p, q \in \mathbb{Z}$ such that $1 = pa_j + q \det G$ and

$$U_j = U_j^{pa_j + q \det G} = (U_j^{a_j})^p (U_j^{\det G})^q \in C^*(\{U_j^{k_j}\}_{j=1}^d, S)$$

for each $j = 1, ..., d$. We have now shown that the generators $\{U_j\}_{j=1}^d$ and S are in $C^*(\{U_j^{k_j}\}_{j=1}^d, S)$. Therefore,

$$\mathcal{O}_{F,G}(\mathbb{T}^d) \subseteq C^*(\{U_j^{k_j}\}_{j=1}^d, S) \subseteq \mathcal{O}_{F,G}(\mathbb{T}^d)$$

and hence, we have equality.

□

Proposition 4.3. Let $\{k_j\}_{j=1}^d \subset \mathbb{N}$ and $\gcd(\det F, \det G) = 1$. Then, as a C^*-subalgebra of $\mathcal{O}_{F,G}(\mathbb{T}^d)$, $C^*(\{U_j^{k_j}\}_{j=1}^d, S)$ is a quotient of $\mathcal{O}_{F,G}(\mathbb{T}^d)$.

Proof. If $\gcd(k_j, a_j) \geq 2$ then represent $k_j = p^{\alpha_j} p_j$ and $a_j = p^{\beta_j} q_j$ where p is prime and p_j, q_j have no factors of p. Suppose $\alpha_j > \beta_j$ then $k_j q_j = p^{\alpha_j - \beta_j} p_j a_j$ and

$$U^{p^{\alpha_j - \beta_j} p_j G_j} = S^* S U^{p^{\alpha_j - \beta_j} p_j G_j} = S^* U_j^{k_j q_j} S \in C^*(\{U_j^{k_j}\}_j, S).$$

As in Lemma 4.2, we obtain

$$U_j^{p^{\alpha_j - \beta_j} p_j \det G} \in C^*(\{U_j^{k_j}\}_j, S)$$

and using $\gcd(\det F, \det G) = 1$,

$$U_j^{p^{\alpha_j - \beta_j} p_j} \in C^*(\{U_j^{k_j}\}_j, S).$$

Suppose $\alpha_j \leq \beta_j$ then $k_j p^{\beta_j - \alpha_j} q_j = p_j a_j$, so similarly

$$U_j^{p_j} \in C^*(\{U_j^{k_j}\}_j, S).$$

These cases show us that we may assume $\gcd(k_j, a_j) = 1$ for all $j = 1, ..., d$.

Suppose $\gcd(k_j, a_j) = 1$ for all $j = 1, ..., d$, then

$$\{(\lambda_1 k_1 \mod a_1, ..., \lambda_d k_d \mod a_d) \,\big|\, \lambda_j \in \{0, ..., a_j - 1\}\} = \oplus_{j=1}^{d} \mathbb{Z}_{a_j}.$$

Hence, given $\lambda \in \mathfrak{I}(F)$, there exists $l_\lambda \in \mathfrak{I}(F)$ and $p_\lambda \in \mathbb{Z}^d$ such that

$$\lambda k = l_\lambda + \mu p_\lambda \qquad \text{(that is, } \lambda_j k_j = l_j + a_j p_j\text{)}$$

where $\mu = (a_1, ..., a_d)$. Set

$$\tilde{S}_\lambda = U^{\lambda k} S \in C^*(\{U_j^{k_j}\}_j, S).$$

Then

$$\begin{aligned}\sum_\lambda \tilde{S}_\lambda \tilde{S}_\lambda^* &= \sum_\lambda (U^{\lambda k} S)(U^{\lambda k} S)^* \\ &= \sum_\lambda U^{l_\lambda} S S^* U^{-l_\lambda} \\ &= \sum_{l_\lambda \in \mathfrak{I}(F)} S_{l_\lambda} S_{l_\lambda}^* = 1\end{aligned}$$

Furthermore, by setting W_j to be the full spectrum unitary $U_j^{k_j}$, we obtain

(1) $\tilde{S}_\nu = W^\nu \tilde{S}$,
(2) $\tilde{S}_\nu^* \tilde{S}_{\nu'} = \delta_\nu^{\nu'}$,
(3) $W_j^{a_j} \tilde{S} = \tilde{S} W^{G_j}$, and
(4) $\sum_\nu W^\nu \tilde{S} \tilde{S}^* W^{-\nu} = 1$.

That is, noting $S = \tilde{S}_{(0,...,0)}$, $C^*(\{U_j^{k_j}\}_{j=1}^d, S)$ is the universal C^*-algebra generated by commuting full spectrum unitaries W_j and isometries $\{\tilde{S}_\nu\}_{\nu \in \mathfrak{I}(F)}$ such that

(1) $\tilde{S}_\nu = W^\nu \tilde{S}$,
(2) $\tilde{S}_\nu^* \tilde{S}_{\nu'} = \delta_\nu^{\nu'}$,
(3) $W_j^{a_j} \tilde{S} = \tilde{S} W^{G_j}$, and
(4) $\sum_\nu W^\nu \tilde{S} \tilde{S}^* W^{-\nu} = 1$.

By Theorem 3.23, the universality of $\mathcal{O}_{F,G}(\mathbb{T}^d)$ implies that there exists a surjective homomorphism of $\mathcal{O}_{F,G}(\mathbb{T}^d)$ onto $C^*(\{U_j^{k_j}\}_{j=1}^d, S)$.

□

5. SPATIAL STRUCTURE OF $\mathcal{O}_{F,G}(\mathbb{T}^d)$

We now turn our attention to a colimit structure of $\mathcal{O}_{F,G}(\mathbb{T}^d)$.

Lemma 5.1. Let $\mathcal{O} = \overline{\operatorname{span}}\{S_\alpha U^\nu S_\beta^* \,\big|\, \alpha = (\alpha_j)_{j=1}^{k_1}, \beta = (\beta_j)_{j=1}^{k_2}, k_1, k_2 \in \mathbb{N}, \alpha_j, \beta_j \in \mathfrak{I}(F), \nu \in \mathbb{Z}^d\}$ where $S_\alpha = S_{\alpha_1} \cdots S_{\alpha_{k_1}}$. Then

$$\mathcal{O} = \mathcal{O}_{F,G}(\mathbb{T}^d)$$

Proof. Begin by noting $S_{\alpha_i} = U^{\alpha_i} S$. With the notation $\alpha_i = (\alpha_{ij})_{j=1}^d \in \mathfrak{I}(F)$

$$U_j S_{\alpha_i} = U^{\alpha_i + e_j} S = \begin{cases} S_{\alpha_i + e_j} & \text{if } \alpha_{ij} < a_j - 1 \\ S_{\alpha_i - (a_j - 1) e_j} U^{G_j} & \text{if } \alpha_{ij} = a_j - 1 \end{cases}$$

where e_j is the vector with 1 in the j-th coordinate and 0 elsewhere. Also,

1. $1 = \sum_\nu S_\nu S_\nu^* \in \mathcal{O}$,
2. $U_j = \sum_\nu S_{\nu+e_j} S_\nu^* \in \mathcal{O}$,
3. $S = \sum_\nu S S_\nu S_\nu^* \in \mathcal{O}$, and
4. $S^* = \sum_\nu S_\nu S_\nu^* S^* \in \mathcal{O}$.

Finally,

$$U_j^* S = U_j^{a_j-1} S U^{-G_j} = S_{(a_j-1)e_j} U^{-G_j};$$

hence, any word in $\{U_j, U_j^*\}_j, S$ and S^* may be expressed as the sum of elements in $\mathcal{O}$.

□

Given $\alpha = (\alpha_j)_{j=1}^k$ where $\alpha_j \in \mathfrak{I}(F)$ and k, the "length of α," is denoted by $|\alpha|$. Let $\gamma : \mathbb{T} \to \operatorname{Aut}(\mathcal{O}_{F,G}(\mathbb{T}^d))$ be the *gauge action* (see [56]) defined by $t \mapsto \gamma_t$ where

$$\gamma_t(S) = tS \qquad \gamma_t(U_j) = U_j \text{ for each } j = 1, ..., d.$$

If $S_\alpha U^\nu S_\beta^* \in \mathcal{O}_{F,G}(\mathbb{T}^d)^\gamma$ (the *fixed point algebra*) then

$$S_\alpha U^\nu S_\beta^* = \gamma_t(S_\alpha U^\nu S_\beta^*) = t^{|\alpha|-|\beta|} S_\alpha U^\nu S_\beta^*$$

implies $|\alpha| = |\beta|$ (since $S_\alpha U^\nu S_\beta^* \neq 0$.) Thus,

$$\mathcal{O}_{F,G}(\mathbb{T}^d)^\gamma \supseteq \overline{\operatorname{span}}\{S_\alpha U^\nu S_\beta^* \,\big|\, |\alpha| = |\beta|, \nu \in \mathbb{Z}^d\}.$$

Furthermore, since $\mathbb{T}$ is compact, averaging over γ with respect to normalized Haar measure produces an expectation of $\mathcal{O}_{F,G}(\mathbb{T}^d)$ onto $\mathcal{O}_{F,G}(\mathbb{T}^d)^\gamma$. Let $E : \mathcal{O}_{F,G}(\mathbb{T}^d) \to \mathcal{O}_{F,G}(\mathbb{T}^d)^\gamma$ be the expectation onto the fixed point algebra $\mathcal{O}_{F,G}(\mathbb{T}^d)^\gamma$ defined by

$$E(x) = \int_{\mathbb{T}} \gamma_t(x)\, dt$$

for each $x \in \mathcal{O}_{F,G}(\mathbb{T}^d)$ (see [56, Section 4.1]). Then for $S_\alpha U^\nu S_\beta^* \in \mathcal{O}_{F,G}(\mathbb{T}^d)$ where $\alpha \in \mathfrak{I}(F)^{k_1}$, $\beta \in \mathfrak{I}(F)^{k_2}$ and $\nu \in \mathbb{Z}^d$ as in Lemma 5.1, note

$$\begin{aligned} E(S_\alpha U^\nu S_\beta^*) &= \int_{\mathbb{T}} t^{|\alpha|-|\beta|} S_\alpha U^\nu S_\beta \, dt \\ &= \frac{S_\alpha U^\nu S_\beta^*}{2\pi} \int_0^{2\pi} e^{2\pi i[|\alpha|-|\beta|]z} dz \\ &= \begin{cases} \frac{S_\alpha U^\nu S_\beta^*}{2\pi} [\frac{e^{2\pi i(|\alpha|-|\beta|)z}}{2\pi i(|\alpha|-|\beta|)}]_0^{2\pi} & \text{if } |\alpha| \neq |\beta| \\ \frac{S_\alpha U^\nu S_\beta^*}{2\pi} [z]_0^{2\pi} & \text{if } |\alpha| = |\beta| \end{cases} \\ &= \delta_{|\alpha|}^{|\beta|} S_\alpha U^\nu S_\beta^* \end{aligned}$$

and by Lemma 5.1,

$$\mathcal{O}_{F,G}(\mathbb{T}^d)^\gamma = E(\mathcal{O}_{F,G}(\mathbb{T}^d)) = \overline{\operatorname{span}}\{S_\alpha U^\nu S_\beta^* \,\big|\, |\alpha| = |\beta|, \nu \in \mathbb{Z}^d\}.$$

Proposition 5.2. Let $\mathcal{A}_k = \overline{\operatorname{span}}\{S_\alpha U^\nu S_\beta^* \,\big|\, |\alpha| = |\beta| = k, \nu \in \mathbb{Z}^d\}$ for $k \in \mathbb{N} \cup \{0\}$. Then $\mathcal{A}_k$ is a unital $*$-subalgebra of $\mathcal{O}_{F,G}(\mathbb{T}^d)^\gamma$ with $\mathcal{A}_k \subset \mathcal{A}_{k+1}$. Moreover,

$$\mathcal{O}_{F,G}(\mathbb{T}^d)^\gamma = \overline{\cup_k \mathcal{A}_k}.$$

Proof. Note $S_\alpha^* S_\beta = \delta_\alpha^\beta$, thus $\mathcal{A}_k$ is closed under multiplication. Furthermore,

$$\begin{aligned} S_\alpha U^\nu S_\beta^* &= S_\alpha \sum_\mu S_\mu S_\mu^* U^\nu S_\beta^* \\ &= \sum_\mu S_{(\alpha,\mu)} U^{\nu_\mu} S_{(\beta,\varpi_\mu)}^* \in \mathcal{A}_{k+1} \end{aligned}$$

where $(\alpha, \mu) = (\alpha_1, ..., \alpha_k, \mu)$ and ϖ_μ and ν_μ are the appropriate vectors with

$$S_\mu^* U^\nu = S^* U^{\nu-\mu} = U^{\nu_\mu} S^* U^{-\varpi_\mu} = U^{\nu_\mu} S_{\varpi_\mu}^*.$$

Finally, $1 = \sum_\nu S_\nu S_\nu^* \in \mathcal{A}_1 \subset \mathcal{A}_k$.

$\square$

Remark 5.3. This result comes as little surprise by Proposition 5.7 of [38] which states that, in general, the fixed point algebra coincides with an inductive limit. This aside, our result leads us to something more interesting.

Lemma 5.4. Let $E_{\alpha\beta} = S_\alpha S_\beta^*$ where $\alpha, \beta \in \mathfrak{I}(F)^k$. Then $\{E_{\alpha\beta}\}_{\alpha,\beta}$ is a system of matrix units for $M_{N^k}(\mathbb{C})$ in $\mathcal{A}_k$.

Proof. We need only check the properties of matrix units:

$$(1) E_{\alpha\beta} E_{\alpha'\beta'} = S_\alpha S_\beta^* S_{\alpha'} S_{\beta'}^* = \delta_\beta^{\alpha'} S_\alpha S_{\beta'}^* = \delta_\beta^{\alpha'} E_{\alpha\beta'},$$

$$\begin{aligned} (2) \sum_\alpha E_{\alpha\alpha} &= \sum_\alpha S_\alpha S_\alpha^* \\ &= \sum_{\alpha_1,\ldots,\alpha_k} S_{\alpha_1} \cdots S_{\alpha_{k-1}} S_{\alpha_k} S_{\alpha_k}^* S_{\alpha_{k-1}}^* \cdots S_{\alpha_1}^* \\ &= \sum_{\alpha_1,\ldots,\alpha_{k-1}} S_{\alpha_1} \cdots S_{\alpha_{k-1}} \Big[\sum_{\alpha_k} S_{\alpha_k} S_{\alpha_k}^*\Big] S_{\alpha_{k-1}}^* \cdots S_{\alpha_1}^* \\ &= \sum_{\alpha_1,\ldots,\alpha_{k-1}} S_{\alpha_1} \cdots S_{\alpha_{k-1}} [1] S_{\alpha_{k-1}}^* \cdots S_{\alpha_1}^* \\ &= ... \\ &= \sum_{\alpha_1} S_{\alpha_1} S_{\alpha_1}^* = 1. \end{aligned}$$

$\square$

Lemma 5.5. [36, Lemma 11.4.8] If $\mathcal{M}$ is a type I factor and B is a C^*-subalgebra of $\mathcal{M}'$, the commutant of $\mathcal{M}$, and A is the C^*-algebra generated by $\mathcal{M} \cup B$, then there is an isomorphism $\varphi : \mathcal{M} \otimes B \to A$ such that $\varphi(m \otimes b) = mb$ for elementary tensors $m \otimes b \in \mathcal{M} \otimes B$.

Note $\mathcal{M}_k := \operatorname{span}\{S_\alpha S_\beta^* \,\big|\, \alpha, \beta \in \mathfrak{I}(F)^k\} \cong M_{N^k}(\mathbb{C})$ (by Lemma 5.4) is a type I factor. Let $B_k = C^*(\{U_j^{a_j^k}\}_{j=1}^d) \cong C(\mathbb{T}^d)$ (since $U_j^{a_j^k}$ is a full spectrum unitary (see Remark 3.22)), then $\mathcal{A}_k$ is the C^*-algebra generated by $\mathcal{M}_k$ and B_k. Indeed, note

$$U_j^{a_j^k} S_\alpha S_\beta^* = S_\alpha U^{(G^k)_j} S_\beta^*$$

for each $\alpha, \beta \in \mathfrak{I}(F)^k$ and $j = 1, ..., d$. Note, in the proof of Lemma 4.1, there exists $\{q_{jl}\}_{j,l=1}^d \subset \mathbb{Z}$ such that

$$\sum_{l=1}^{d} q_{jl}(G^k)_l t = (\det G^k) t_j$$

for each $j = 1, ..., d$. Hence, $U_j^{\det G^k} \in C^*(\{U^{(G^k)_j}\}_{j=1}^d)$. Further note $\{U_j^{\det G^k}\}_{j=1}^d$ generates the faithful image of $C(\mathbb{T}^d)$ in $\mathcal{O}_{F,G}(\mathbb{T}^d)$ (since $\det G^k = (\det G)^k \neq 0$ and the left action is injective.) Thus, $\mathcal{A}_k$ is the C^*-algebra generated by $\mathcal{M}_k$ and B_k. Furthermore,

$$U_j^{a_j^k} S_\alpha S_\beta^* = S_\alpha U^{(G^k)_j} S_\beta^* = S_\alpha (S_\beta U^{-(G^k)_j})^* = S_\alpha (U_j^{-a_j^k} S_\beta)^* = S_\alpha S_\beta^* U_j^{a_j^k}$$

for each $j = 1, ..., d$ and $\alpha, \beta \in \mathfrak{I}(F)^k$; hence, the generators of B commute with the generators of $\mathcal{M}$. Thus, B may be viewed as a C^*-subalgebra of $\mathcal{M}'$. Hence, we use the previous lemma.

Theorem 5.6.

$$\mathcal{A}_k \cong C(\mathbb{T}^d) \otimes M_{N^k}(\mathbb{C})$$

Proof. By Lemma 5.5, we obtain an isomorphism $\psi_k : B_k \otimes \mathcal{M}_k \to \mathcal{A}_k$. Let $\psi_{B_k} : B_k \to C(\mathbb{T}^d)$ and $\psi_{\mathcal{M}_k} : \mathcal{M}_k \to M_{N^k}(\mathbb{C})$ be isomorphisms, then

$$(\psi_{B_k} \otimes \psi_{\mathcal{M}_k}) \circ (\psi_k)^{-1} : \mathcal{A}_k \to C(\mathbb{T}^d) \otimes M_{N^k}(\mathbb{C})$$

is an isomorphism.

□

Hence, for the maps $\varphi_k : C(\mathbb{T}^d) \otimes M_{N^k}(\mathbb{C}) \cong \mathcal{A}_k \hookrightarrow \mathcal{A}_{k+1} \cong C(\mathbb{T}^d) \otimes M_{N^{k+1}}(\mathbb{C})$, we have

$$\mathcal{O}_{F,G}(\mathbb{T}^d)^\gamma = \operatorname{colim}(M_{N^k}(C(\mathbb{T}^d)), \varphi_k).$$

We use the identification $C(\mathbb{T}^d) \otimes M_{N^k}(\mathbb{C}) \cong M_{N^k}(C(\mathbb{T}^d))$.

We now define the notion of a crossed product by an endomorphism as discussed in [62].

Definition 5.7. [62] A *crossed product* (of multiplicity 1) for an endomorphism ρ of a unital C^*-algebra A with

$$A_\infty = \operatorname{colim}_{k \in \mathbb{N}}(A, \rho) \neq 0$$

is the (unique) unital C^*-algebra $A \rtimes_\rho \mathbb{N}$ together with a homomorphism $\iota : A \to A \rtimes_\rho \mathbb{N}$ with $\iota(1_A) = 1_{A \rtimes_\rho \mathbb{N}}$, and an isometry $s \in A \rtimes_\rho \mathbb{N}$ such that

1. $\iota(\rho(a)) = s\iota(a)s^*$ for all $a \in A$
2. for every representation π of A and isometry t satisfying $\pi(\rho(a)) = t\pi(a)t^*$ for all $a \in A$, there exists a non-degenerate representation $\pi \times T$ of $A \rtimes_\rho \mathbb{N}$ with
$$(\pi \times T) \circ \iota = \pi \qquad \text{and} \qquad (\pi \times T)(s) = t.$$
3. $A \rtimes_\rho \mathbb{N}$ is generated by $\iota(A)$ and s.

Let $\rho_k : C(\mathbb{T}^d) \otimes M_{N^k}(\mathbb{C}) \to C(\mathbb{T}^d) \otimes M_{N^{k+1}}(\mathbb{C})$ be defined on basic tensors of $C(\mathbb{T}^d) \otimes M_{N^k}(\mathbb{C})$ by

$$f \otimes E_{\alpha\beta} \mapsto f \otimes E_{(0,\alpha)(0,\beta)}$$

where $\alpha, \beta \in \mathfrak{I}(F)^k$ and $f \in C(\mathbb{T}^d)$. Note that since $\{E_{\alpha\beta}\}_{\alpha,\beta\in\mathfrak{I}(F)^k}$ is linearly independent, the definition of ρ_k extends, by linearity, to all of $C(\mathbb{T}^d) \otimes M_{N^k}(\mathbb{C})$.

Lemma 5.8. Let $\rho_k : C(\mathbb{T}^d) \otimes M_{N^k}(\mathbb{C}) \to C(\mathbb{T}^d) \otimes M_{N^{k+1}}(\mathbb{C})$ be defined by

$$f \otimes E_{\alpha\beta} \mapsto f \otimes E_{(0,\alpha)(0,\beta)}$$

on the basic tensors, $f \otimes E_{\alpha\beta}$, of $C(\mathbb{T}^d) \otimes M_{N^k}(\mathbb{C})$ where $\alpha, \beta \in \mathfrak{I}(F)^k$ and $f \in C(\mathbb{T}^d)$. Then ρ_k is a (continuous) $*$-homomorphism for each $k \in \mathbb{N}$ and the following diagram commutes

$$\begin{tikzcd}
\cdots \arrow[r, "\varphi_{k-1}"] & \mathcal{C}(\mathbb{T}^d) \otimes M_{N^k}(\mathbb{C}) \arrow[r, "\varphi_k"] & \mathcal{C}(\mathbb{T}^d) \otimes M_{N^{k+1}}(\mathbb{C}) \arrow[r, "\varphi_{k+1}"] & \cdots \\
\cdots \arrow[r, "\varphi_{k-1}"] \arrow[ur, "\rho_{k-1}"] & \mathcal{C}(\mathbb{T}^d) \otimes M_{N^k}(\mathbb{C}) \arrow[r, "\varphi_k"] \arrow[ur, "\rho_k"] & \mathcal{C}(\mathbb{T}^d) \otimes M_{N^{k+1}}(\mathbb{C}) \arrow[r, "\varphi_{k+1}"] \arrow[ur, "\rho_{k+1}"] & \cdots
\end{tikzcd}$$

Proof. Note for any $\sum_{\alpha,\beta\in\mathfrak{I}(F)^k} f_{\alpha,\beta} \otimes E_{\alpha\beta}$ and $\sum_{\alpha',\beta'\in\mathfrak{I}(F)^k} f_{\alpha',\beta'} \otimes E_{\alpha'\beta'}$

$$\begin{aligned}
\rho_k\Big(\Big(\sum_{\alpha,\beta\in\mathfrak{I}(F)^k} & f_{\alpha,\beta} \otimes E_{\alpha\beta}\Big)\Big(\sum_{\alpha',\beta'\in\mathfrak{I}(F)^k} f_{\alpha',\beta'} \otimes E_{\alpha'\beta'}\Big)\Big) \\
&= \rho_k\Big(\sum_{\alpha,\beta,\alpha',\beta'\in\mathfrak{I}(F)^k} f_{\alpha,\beta} f_{\alpha',\beta'} \otimes E_{\alpha\beta}E_{\alpha'\beta'}\Big) \\
&= \rho_k\Big(\sum_{\alpha,\beta'\in\mathfrak{I}(F)^k} f_{\alpha,\beta} f_{\beta,\beta'} \otimes E_{\alpha\beta'}\Big) \\
&= \sum_{\alpha,\beta'\in\mathfrak{I}(F)^k} f_{\alpha,\beta} f_{\beta,\beta'} \otimes E_{(0,\alpha)(0,\beta')}\Big) \\
&= \Big(\sum_{\alpha,\beta\in\mathfrak{I}(F)^k} f_{\alpha,\beta} \otimes E_{(0,\alpha)(0,\beta)}\Big)\Big(\sum_{\alpha',\beta'\in\mathfrak{I}(F)^k} f_{\alpha',\beta'} \otimes E_{(0,\alpha')(0,\beta')}\Big) \\
&= \rho_k\Big(\sum_{\alpha,\beta\in\mathfrak{I}(F)^k} f_{\alpha,\beta} \otimes E_{\alpha\beta}\Big)\rho_k\Big(\sum_{\alpha',\beta'\in\mathfrak{I}(F)^k} f_{\alpha',\beta'} \otimes E_{\alpha'\beta'}\Big)
\end{aligned}$$

Recall that $\mathcal{A}_k \subset \mathcal{A}_{k+1}$. Let i_k denote the inclusion and recall, for any $S_\alpha U^\nu S_\beta^* \in \mathcal{A}_k$

$$i_k(S_\alpha U^\nu S_\beta^*) = S_\alpha\Big(\sum_{\mu\in\mathfrak{I}(F)} S_\mu S_\mu^*\Big)U^\nu S_\beta^* = \sum_{\mu\in\mathfrak{I}(F)} S_{(\alpha,\mu)} U^{\nu_\mu} S^*_{(\beta,\varpi_\mu)}$$

for appropriate $\nu_\mu \in \mathbb{Z}^d$ and $\varpi_\mu \in \mathfrak{I}(F)$ as in Proposition 5.2. Thus for φ_k defined in the paragraph proceeding Theorem 5.6,

$$\varphi_k(U^\nu \otimes E_{\alpha\beta}) = \sum_{\mu\in\mathfrak{I}(F)} U^{\nu_\mu} \otimes E_{(\alpha,\mu)(\beta,\varpi_\mu)}$$

for the appropriate $\nu_\mu \in \mathbb{Z}^d$ and $\varpi_\mu \in \mathfrak{I}(F)$. Furthermore, since

$$\begin{aligned} i_{k+1}(S_{(0,\alpha)}U^\nu S^*_{(0,\beta)}) &= SS_\alpha(\sum_{\mu\in\mathfrak{I}(F)} S_\mu S^*_\mu)U^\nu S^*_\beta S^* \\ &= S(\sum_{\mu\in\mathfrak{I}(F)} S_{(\alpha,\mu)}U^{\nu_\mu} S^*_{(\beta,\varpi_\mu)})S^* \\ &= \sum_{\mu\in\mathfrak{I}(F)} S_{(0,\alpha,\mu)}U^{\nu_\mu} S^*_{(0,\beta,\varpi_\mu)} \end{aligned}$$

we obtain

$$\varphi_{k+1}(U^\nu \otimes E_{(0,\alpha)(0,\beta)}) = \sum_{\mu\in\mathfrak{I}(F)} U^{\nu_\mu} \otimes E_{(0,\alpha,\mu)(0,\beta,\varpi_\mu)}.$$

Hence,

$$\begin{aligned} (\rho_{k+1}\circ\varphi_k)(U^\nu \otimes E_{\alpha\beta}) &= \rho_{k+1}(\sum_{\mu\in\mathfrak{I}(F)} U^{\nu_\mu} \otimes E_{(\alpha,\mu)(\beta,\varpi_\mu)}) \\ &= \sum_{\mu\in\mathfrak{I}(F)} U^{\nu_\mu} \otimes E_{(0,\alpha,\mu)(0,\beta,\varpi_\mu)} \end{aligned}$$

and

$$\begin{aligned} (\varphi_{k+1}\circ\rho_k)(U^\nu \otimes E_{\alpha\beta}) &= \varphi_{k+1}(U^\nu \otimes E_{(0,\alpha)(0,\beta)}) \\ &= \sum_{\mu\in\mathfrak{I}(F)} U^{\nu_\mu} \otimes E_{(0,\alpha,\mu)(0,\beta,\varpi_\mu)} \\ &= (\rho_{k+1}\circ\varphi_k)(U^\nu \otimes E_{\alpha\beta}) \end{aligned}$$

for all $U^\nu \otimes E_{\alpha\beta} \in C(\mathbb{T}^d) \otimes M_{N^k}(\mathbb{C})$. The result now follows from the linearity and continuity of ρ_k and φ_k.

□

Let $j_k : \mathcal{A}_k \hookrightarrow \mathcal{O}_{F,G}(\mathbb{T}^d)^\gamma$ be the natural inclusion $*$-homomorphism for each $k \in \mathbb{N}$ and let $\tilde{\rho}_k : \mathcal{A}_k \to \mathcal{O}_{F,G}(\mathbb{T}^d)^\gamma$ be the composition $j_{k+1} \circ \Psi^{-1}_{k+1} \circ \rho_k \circ \Psi_k$ for $k \in \mathbb{N}$ where Ψ_k is the isomorphism in Theorem 5.6. Then the solid diagram

$$\begin{tikzcd}
 & & \mathcal{O}_{F,G}(\mathbb{T}^d)^\gamma & & & & \\
\cdots \arrow[r, "i_{k-2}"'] & \mathcal{A}_{k-1} \arrow[ur, "\tilde{\rho}_{k-1}"] \arrow[r, "i_{k-1}"'] & \mathcal{A}_k \arrow[u, "\tilde{\rho}_k"'] \arrow[r, "i_k"'] & \mathcal{A}_{k+1} \arrow[ul, "\tilde{\rho}_{k+1}"'] \arrow[r, "i_{k+1}"'] & {} & \cdots \arrow[r] & \mathcal{O}_{F,G}(\mathbb{T}^d)^\gamma \arrow[ullll, dashed, "\rho"']
\end{tikzcd}$$

commutes and hence, there exists a (unique) $*$-endomorphism ρ on $\mathcal{O}_{F,G}(\mathbb{T}^d)^\gamma$ making the entire diagram commute. A diagram chase ensures that

$$\rho(S_\alpha f S^*_\beta) = S_{(0,\alpha)} f S^*_{(0,\beta)}$$

for $S_\alpha f S^*_\beta \in \mathcal{O}_{F,G}(\mathbb{T}^d)^\gamma$.

Theorem 5.9.

$$\mathcal{O}_{F,G}(\mathbb{T}^d) \cong \mathcal{O}_{F,G}(\mathbb{T}^d)^\gamma \rtimes_\rho \mathbb{N}$$

where $\rho \in \operatorname{End}(\mathcal{O}_{F,G}(\mathbb{T}^d)^\gamma)$ is defined above.

Proof. Let $(\iota, s) : \mathcal{O}_{F,G}(\mathbb{T}^d)^\gamma \to \mathcal{O}_{F,G}(\mathbb{T}^d)^\gamma \rtimes_\rho \mathbb{N}$ be the universal representation as described in Definition 5.7. Then let $\iota_0 : \mathcal{O}_{F,G}(\mathbb{T}^d)^\gamma \to \mathcal{O}_{F,G}(\mathbb{T}^d)$ be the natural inclusion and let $S \in \mathcal{O}_{F,G}(\mathbb{T}^d)$ denote the isometry$S_{(0,0,\dots,0)}$ defined in Theorem 3.23. Note

$$\iota_0(\rho(S_\alpha U^\nu S_\beta^*)) = S_{(0,\alpha)} U^\nu S_{(0,\beta)}^* = S S_\alpha U^\nu S_\beta^* S^* = S \iota_0(S_\alpha U^\nu S_\beta^*) S^*$$

for all generators $S_\alpha U^\nu S_\beta^* \in \mathcal{O}_{F,G}(\mathbb{T}^d)^\gamma$. Hence,

$$\iota_0(\rho(x)) = S \iota_0(x) S^*$$

for each $x \in \mathcal{O}_{F,G}(\mathbb{T}^d)^\gamma$. Furthermore, note $C^*(\mathcal{O}_{F,G}(\mathbb{T}^d)^\gamma, S) = \mathcal{O}_{F,G}(\mathbb{T}^d)$. Therefore, there exists a homomorphism

$$\tau : \mathcal{O}_{F,G}(\mathbb{T}^d)^\gamma \rtimes_\rho \mathbb{N} \to \mathcal{O}_{F,G}(\mathbb{T}^d)$$

such that

$$\tau \circ \iota = \iota_0 \qquad \text{and} \qquad \tau(s) = S.$$

Recall by Lemma 5.1, there exists $x \in \mathcal{O}_{F,G}(\mathbb{T}^d)^\gamma$ such that $\rho(x^\nu) = U^\nu$ for each $\nu \in \mathfrak{I}(F)$. Then

(1) Given $j \in \{1, \dots, d\}$,

$$\iota(U_j)^{a_j} s = \iota(\rho(x))^{a_j} s = s \iota(x)^{a_j} s^* s = s \iota(x^{a_j})$$

and

$$x^{a_j} = S^* S x^{a_j} S^* S = S^* \rho(x)^{a_j} S = S^* U^{a_j} S = U^{G_j}.$$

Hence,

$$\iota(U_j)^{a_j} s = s \iota(U)^{G_j}$$

for each j.

(2) Given $\nu \in \mathfrak{I}(F)$,

$$s^* \iota(U)^\nu s = s^* \iota(\rho(x))^\nu s = s^* s \iota(x)^\nu s^* s = \iota(x)^\nu$$

and

$$x^\nu = S^* S x^\nu S^* S = S^* \rho(x)^\nu S = S^* U^\nu S = \delta_0^\nu.$$

Hence,

$$s^* \iota(U)^\nu s = \delta_0^\nu$$

for each $\nu \in \mathfrak{I}(F)$.

(3) Finally,

$$\sum_\nu \iota(U)^\nu s s^* \iota(U)^{-\nu} = \sum_\nu \iota(U)^\nu \iota(\rho(1)) \iota(U)^{-\nu} = \sum_\nu \iota(U^\nu S S^* U^{-\nu}) = \iota(1) = 1.$$

Therefore, by Theorem 3.23, there exists a homomorphism

$$\sigma : \mathcal{O}_{F,G}(\mathbb{T}^d) \to \mathcal{O}_{F,G}(\mathbb{T}^d)^\gamma \rtimes_\rho \mathbb{N}$$

such that

$$U_j \mapsto \iota(U_j) \text{ for each } j = 1, ..., d \qquad \text{and} \qquad S \mapsto s.$$

Then

$$(\tau \circ \sigma)(U_j) = (\tau \circ \iota)(U_j) = \iota_0(U_j) = U_j$$

for each $j = 1, ..., d$ and

$$(\tau \circ \sigma)(S) = \tau(s) = S.$$

Hence, $\tau \circ \sigma = \mathrm{id}_{\mathcal{O}_{F,G}(\mathbb{T}^d)}$. Furthermore, for each $S_\alpha U^\nu S_\beta^* \in \mathcal{O}_{F,G}(\mathbb{T}^d)^\gamma$,

$$\begin{aligned}
(\sigma \circ \tau)(\iota(S_\alpha U^\nu S_\beta^*)) &= \sigma(S_\alpha U^\nu S_\beta^*) \\
&= \sigma(U^{\alpha_1} S U^{\alpha_2} S \cdots U^{\alpha_k} S U^\nu S^* U^{-\beta_k} S^* U^{-\beta_{k-1}} \cdots S^* U^{-\beta_1} \\
&= \iota(U^{\alpha_1}) s \iota(U^{\alpha_2}) s \cdots \iota(U^{\alpha_k}) s \iota(U^\nu) s^* \iota(U^{-\beta_k}) s^* \iota(U^{-\beta_{k-1}}) \cdots s^* \iota(U^{-\beta_1}) \\
&= \iota(U^{\alpha_1}) s \iota(U^{\alpha_2}) s \cdots \iota(U^{\alpha_{k-1}}) S^* \iota(U^{\alpha_k} S U^\nu S^* U^{-\beta_k})) s^* \iota(U^{-\beta_{k-1}}) \cdots s^* \iota(U^{-\beta_1}) \\
&= \cdots = \iota(S_\alpha U^\nu S_\beta^*)
\end{aligned}$$

and

$$(\sigma \circ \tau)(s) = \sigma(S) = s.$$

Thus, by Lemma 5.1, $\sigma \circ \tau = \mathrm{id}_{\mathcal{O}_{F,G}(\mathbb{T}^d)^\gamma \rtimes_\rho \mathbb{N}}$ and so,

$$\mathcal{O}_{F,G}(\mathbb{T}^d) \cong \mathcal{O}_{F,G}(\mathbb{T}^d)^\gamma \rtimes_\rho \mathbb{N} \cong (\mathrm{colim}_k(M_{N^k}(C(\mathbb{T}^d))) \rtimes_\rho \mathbb{N}$$

□

6. Acknowledgements

This paper is the first product of my doctoral dissertation at the University of Calgary. It would not have been possible without the guidance and help of Dr. Berndt Brenken and Dr. Marcelo Laca for their insight on corrections and enhancements of the material discussed here.

I am also indebted to NSERC, the Department of Mathematics and Statistics at the University of Calgary and the Department of Mathematics and Statistics at the University of Regina in providing funding to finance my mathematical studies.

References

[1] an Huef, A., and Raeburn, I., *The ideal structure of Cuntz-Krieger algebras*, Ergodic Theory Dyn. Sys. 17 (1997) 611-624.

[2] Bates, T., Hong, J. H., Raeburn, I., and Szymanski, W., *The ideal structure of C∗-algebras of infinite graphs*, Illinois J. Math 46 (2002), 1159-1176.

[3] Bates, T., Pask, D., Raeburn, I., and Szymanski, W., *The C^*-algebras of row-finite graphs*, New York J. Math. 6 (2000), 307-324.

[4] Blackadar, B., K-theory for Operator Algebras, M. S. R. I. Monographs, vol. 5, Springer-Verlag, Berlin and New York, 1986.

[5] Blackadar, B., Operator Algebras: Theory of C*-Algebras and von Neumann Algebras, Encyclopaedia of Mathematical Sciences, vol. 122, Springer-Verlag, Berlin, 2006.

[6] Brenken, B., *The local product structure of expansive automorphisms of solenoids and their associated C^*-algebras*, Can. J. Math., 48, (1996), 692-709.

[7] Brenken, B., *C^*-algebras associated with topological relations*, J. Ramanujan Math. Soc. 19, No. 1 (2004), 1-21.

[8] Brenken, B., *Endomorphisms of type I von Neumann algebras with discrete center*, J. Operator Theory 51 (2004), no. 1, 19-34.

[9] Brenken, B., *The isolated ideal of a correspondence associated with a topological quiver*. New York J. Math., 12, (2006), 1-16.

[10] Brenken, B., *A Dynamical Core for Topological Directed Graphs*, Munster J. of Math. 3 (2010), 111-144.

[11] Brenken, B., *Topological Quivers as Multiplicity Free Relations*, Math. Scand., 106, (2010), 217-242.

[12] Brown, N.P., and Ozawa, N., C^*-Algebras and Finite-Dimensional Approximations, Graduate Studies in Mathematics, 88. Amer. Math. Soc. Providence, Rhode Island, 2008.

[13] Conway, J.B., A Course in Functional Analysis, Second Edition, Graduate Texts in Mathematics, 96. Springer, New York, 1990.

[14] Cuntz, J., *Simple C^*-algebras generated by isometries*. Comm. Math. Phys. 57 (1977), no. 2, 173-185.

[15] Cuntz, J., and Krieger, W., *A class of C^*-algebras and topological Markov chains*, Inventiones Math., 56 (1980), 251-268.

[16] Davidson, K., C^*-Algebras by Example, Fields Institute Monograph, Amer. Math. Soc. Providence, Rhode Island, 1996.

[17] Deaconu, V., *Groupoids associated with endomorphisms*, Trans. Amer. Math. Soc. 347 (1995), 1779-1786.

[18] Deaconu, V., *A path model for circle algebras*, J. Operator Theory 34 (1995), 57-89.

[19] Deaconu, V., *Generalized Cuntz-Krieger algebras*, Proc. Amer. Math. Soc. 124 (1996), 3427-3435.

[20] Deaconu, V., *Continuous graphs and C^*-algebras*, in Operator theoretical methods (Timisoara, 1998), 137-149, Theta Found., Bucharest, 2000.

[21] Deaconu, V., and Muhly, P., *C^*-algebras associated with branched coverings*, Proc. Amer. Math. Soc. 129 (2001), 1077-1086.

[22] Dummit, D.S., and Foote, R.M., Abstract Algebra, 3rd edition, Wiley and Sons, 2004.

[23] Evans, D.E., *The C^*-algebras of topological Markov chains*, Lecture Notes, Tokyo Metropolitan University (1983).

[24] Exel, R., an Huef, A., and Raeburn, I., *Purely Infinite Simple C^*-algebras associated to Integer Dilation Matrices*, Indiana Univ. Math. J. 60 (2011), no. 3, 1033-1058.

[25] Fowler, N.J., Laca, M., and Raeburn, I., *The C^*-algebras of infinite graphs*, Proc. Amer. Math. Soc. 8 (2000), 2319-2327.

[26] Fowler, N.J., Muhly, P.S., and Raeburn, I., *Representations of Cuntz-Pimsner Algerbas*, Indiana Univ. Math. J., 52(3) (2003), 569-605.

[27] Fowler, N.J., Raeburn, I., *The Toeplitz algebra of a Hilbert bimodule*, Indiana Univ. Math. J. 48 (1999), 155-181.

[28] Gabriel, P., *Unzerlegbare Darstellungen I* (Oberwolfach 1970), Manuscr. Math. 6 (1972), 71-103.

[29] Hajac, P. M., Matthes, R., and Szymański, W., *Graph C^*-algebras and $\mathbb{Z}_2$-quotients of quantum spheres*. Proceedings of the XXXIV Symposium on Mathematical Physics (Toru, 2002). Rep. Math. Phys. 51 (2003), no. 2-3, 215-224.

[30] Hajac, P.M., Matthes, R., and Szymański, W., *Quantum real projective space, disc and spheres*, Algebr. Represent. Theory 6 (2003), 169-192.

[31] Hong, J.H., and Szymański, W., *Quantum spheres and projective spaces as graph algebras*, Comm. Math. Phys. 232 (2002), 157-188.

[32] Han, D., Jing, W., Larson, D., and Mohapatra, R., *Riesz bases and their dual modular frames in Hilbert C^*-modules.* J. Math. Anal. Appl. 343 (2008), no. 1, 246-256.

[33] Hatcher, A., Algebraic Topology, Cambridge University Press, New York, 2001.

[34] Ji, R., *On Crossed Product C^*-Algebras Associated with Furstenberg Transformations on Tori*, PhD Thesis, State University of New York, Stony Brook, 1986.

[35] Kadison, R.V., and Ringrose, J.R., Fundamentals of the theory of operator algebras. Vol. I. Elementary theory, Graduate Studies in Mathematics, 15. Amer. Math. Soc., Providence, RI, 1997.

[36] Kadison, R.V., and Ringrose, J.R., Fundamentals of the theory of operator algebras. Vol. II. Advanced theory, Graduate Studies in Mathematics, 16. Amer. Math. Soc., Providence, RI, 1997.

[37] Kajiwara, T., and Watatani, Y., *Hilbert C^*-bimodules and continuous Cuntz-Krieger algebras.* J. Math. Soc. Japan 54 (2002), no. 1, 35-59.

[38] Katsura, T., *On C^*-algebras associated with C^*-correspondences.* J. of Functional Analysis 217 (2004), 366-401.

[39] Katsura, T., *A class of C^*-algebras generalizing both graph algebras and homeomorphism C^*-algebras. I. Fundamental results.* Trans. Amer. Math. Soc. 356 (2004), no. 11, 4287-4322.

[40] Katsura, T., *A class of C^*-algebras generalizing both graph algebras and homeomorphism C^*-algebras. II. Examples.* Internat. J. Math. 17 (2006), no. 7, 791-833.

[41] Katsura, T., *A class of C^*-algebras generalizing both graph algebras and homeomorphism C^*-algebras. III. Ideal structures.* Ergodic Theory Dynam. Systems 26 (2006), no. 6, 1805-1854.

[42] Katsura, T., *A class of C^*-algebras generalizing both graph algebras and homeomorphism C^*-algebras. IV. Pure infiniteness.* J. Funct. Anal. 254 (2008), no. 5, 1161-1187.

[43] Kumjian, A.,*Notes on C^*-algebras of graphs,* Contemporary Math. 228 (1998) 189-200.

[44] Kumjian, A., and Pask, D., *Higher rank graph C^*-algebras*, New York J. Math. 6 (2000), 1-20.

[45] Kumjian, A., and Pask, D., *Actions of $\mathbb{Z}^k$ associated to higher rank graphs*, Ergodic Theory & Dynamical Systems, 23 (2003), 1153-1172.

[46] Kumjian, A., Pask, D., and Raeburn, I., *Cuntz-Krieger algebras of directed graphs*, Pacific J. Math. 184 (1998), 161-174.

[47] Kumjian, A., Pask, D., Raeburn, I., and Renault, J., *Graphs, groupoids, and Cuntz-Krieger algebras*, J. Funct. Anal. 144 (1997), 505-541.

[48] Lance, E.C., Hilbert C^*-modules: A toolkit for operator algebraists, London Mathematical Society Lecture Note Series, vol. 210, Cambridge University Press, 1995.

[49] Mann, M.H., Raeburn, I., and Sutherland, C.E., *Representations of finite groups and Cuntz-Krieger algebras*, Bull. Austral. Math. Soc. 46 (1992), 225243.

[50] Mann, M.H., Raeburn, I., and Sutherland, C.E., *Representations of compact groups, Cuntz-Krieger algebras, and groupoid C^*-algebras* in Miniconference on probability and analysis (Sydney, 1991), 135144, Proc. Centre Math. Appl. Austral. Nat. Univ., 29, Austral. Nat. Univ., Canberra, 1992.

[51] McCann, S.J., *C^*-algebras associated with topological group quivers*, PhD Thesis, University of Calgary, Calgary, Alberta, Canada, 2012.

[52] McCann, S. J., *C^*-algebras associated with topological group quivers I: generators, relations and spatial structure*, preprint

[53] Muhly, P., and Solel, B., *Tensor algebras over C^*-correspondences (representations, dilations, and C^*-envelopes)*, J. Funct. Anal. 158 (1998), 389-457.

[54] Muhly, P., and Solel, B., *On the Morita Equivalence of Tensor algebras*, Proc. London Math. Soc. 81 (2000), 113-168.

[55] Muhly, P., and Tomforde, M., *Adding tails to C^*-correspondences*, Doc. Math. 9 (2004), 79-106.

[56] Muhly, P., and Tomforde, M., *Topological quivers.* Internat. J. Math. 16 (2005), no. 7, 693-755.

[57] Munkres, J. R., Topology, 2nd edition, Prentice-Hall, 2000.

[58] Pask, D., and Sutherland, C.E., *Filtered inclusions of path algebras; a combinatorial approach to Doplicher-Roberts duality*, J. Operator Theory 31 (1994), 99121.

[59] Paulsen, V.I., Completely bounded maps and operator algebras, Cambridge Studies in Advanced Math., 78, Cambridge University Press, Cambridge, 2002.

[60] Pimsner, M., *A class of C^*-algebras generating both Cuntz-Krieger algebras and crossed products by $\mathbb{Z}$*, in Free Probability Theory, fields inst. Commun., vol. 12, Amer. Math. Soc., Providence, 1997, pages 189-212.

[61] Schweizer, J., *Crossed Product by C^*-correspondences and Cuntz-Pimsner Algebras*, 'in C^*-algebras, Proceedings of the SFB-Workshop on C^*-algebras, Muenster, 1999,' (Eds.) J. Cuntz, S. Echterhoff, Springer Verlag, Berlin, 2000.

[62] Stacey, P.J., *Crossed products of C^*-algebras by endomorphisms*, J. Austral. Math. Soc. (Series A) 54 (1993), 204-212.

[63] Raeburn, I., Williams, D.P., Morita Equivalence and Continuous-Trace C^*-Algebras, Math. Surveys & Monographs, vol. 60, Amer. Math. Soc., Providence, 1998.

[64] Rieffel, M., *C^*-algebras associated with irrational rotations*, Pacific J. Math. 93(2) (1981), 415-429.

[65] Rϕrdam, M., Larsen, F., Lausten, N.J., An Introduction to K-theory for C*-algebras. 256 pp. London Mathematical Society, Student Text 49, Cambridge University Press, Cambridge, 2000.

[66] Rϕrdam, M., Stϕrmer, E., Operator Algebras and Non-Commutative Geometry, Vol VII: Classification of Nuclear C*-Algebras. Entropy in Operator Algebras. Encyclopaedia of Mathematical Sciences 126. Springer Verlag, Heidelberg, 2001.

[67] Walters, P., An introduction to ergodic theory, Springer, New York, 1982.

[68] Yamashita, S., *Circle Correspondence C^*-algebras*, Houston J. Math. 37 (2011), no. 4, 1181-1202.

DEPARTMENT OF MATHEMATICS AND STATISTICS, UNIVERSITY OF REGINA, REGINA, SK, S4S 0A2
E-mail address: `mccann1s@uregina.ca`